

PLATE XIV.

Fig. 1. One of the superior ventral bristles (with whorls of spikes) of <i>Leanira japanensis</i> , . . . . .	× 250.
Fig. 2. One of the upper and larger ordinary ventral bristles of the same, . . . . .	× 250.
Fig. 3. Ventral bristle of <i>Leanira laevis</i> , . . . . .	× 250.
Fig. 4. Serrated bristle, from the inferior division of the tenth foot of <i>Eulepis wyvillei</i> , . . . . .	× 400.
Fig. 5. One of the powerful golden dorsal hooks, from the middle region of the body in the same species, . . . . .	× 100.
Fig. 6. One of the strong winged bristles, from the inferior division of the foot, . . . . .	× 110.
Fig. 7. Dorsal hook of <i>Eulepis challengeriae</i> , . . . . .	× 400.
Fig. 8. Solitary pectinate bristle, from the superior edge of the inferior division of the foot of the same, . . . . .	× 600.
Fig. 9. One of the short, transversely barred bristles of <i>Nephthys trisso-phylloides</i> , . . . . .	× 210.
Fig. 10. One of the superior elongated bristles of the same, . . . . .	× 90.
Fig. 11. A long posterior bristle from another (smaller) example, having the parasitic Infusorian attached, . . . . .	× 90.
Fig. 12. Barred bristle of <i>Nephthys phyllobranchia</i> , . . . . .	× 400.
Fig. 13. One of the elongated bristles of the same, . . . . .	× 90.
Fig. 14. Bristle (average) in profile, from the anterior third of <i>Phyllocoete (Anaitis) sanctæ-vincentis</i> , . . . . .	× 800.
Fig. 15. Antero-posterior view of one of the same, . . . . .	× 800.
Figs. 16, 17. Two varieties of bristles, from the anterior third of <i>Eulalia capensis</i> , . . . . .	× 530.