

PLATE XV_A.

Fig. 1.	Bristle of <i>Phyllodoce duplex</i> ,	×	800.
Fig. 2.	Bristle of <i>Genetyllis</i> (?) <i>oculata</i> .	Viewed somewhat antero-						×	350.
		posteriorly,		
Fig. 3.	Another of the same seen laterally,	×	350.
Fig. 4.	Bristle of <i>Greeffia oahuensis</i> ,	×	430.
Fig. 5.	Dorsal bristle of <i>Dalhousia atlantica</i> ,	×	50.
Fig. 6.	One of the longer (superior) ventral bristles of the same,	×	90.
Fig. 7.	One of the forms with shorter tips,	×	210.
Fig. 8.	One of the posterior bristles (with short tip) of a fragmentary								
	<i>Hesione</i> , from St. Thomas's, West Indies,	×	400.
Fig. 9.	Another with a longer extremity,	×	400.
Fig. 10.	Average bristle of <i>Hesione</i> (<i>Fallacia</i>) <i>pantherina</i> ,	×	400.
Fig. 11.	Simple solitary bristle of <i>Salvatoria kerguelensis</i> ,	×	400.
Fig. 12.	Jointed inferior bristle of the same,	×	400.
Fig. 13.	One of the longer bristles of <i>Eusyllis kerguelensis</i> ,	×	400.
Fig. 14.	Bristle of <i>Syllis gigantea</i> ,	×	250.
Fig. 15.	Bristle of <i>Autolytus maclearanus</i> ,	×	400.
Fig. 16.	Setigerous process of one of the feet, with the bristles of <i>Syllis</i>								
	<i>setubalensis</i> ,	×	400.
Fig. 17.	Setigerous process of a posterior foot of the foregoing, with the								
	bristles,	×	400.
Fig. 18.	Setigerous lobe of the anterior region of an attached female bud of								
	<i>Syllis ramosa</i> (13 on the Plate by mistake),	×	400.
Fig. 19.	Tip of one of the broad flattened translucent bristles in a free								
	female bud of the same,	×	400.
Fig. 20.	Setigerous region of one of the anterior feet of <i>Syllis brasiliensis</i> ,	×	400.
Fig. 21.	Upper bristle of <i>Syllis capensis</i> ,	×	400.
Fig. 22.	Bristle of <i>Sphaerosyllis kerguelensis</i> ,	×	400.