

PLATE XVIIA.

Fig. 1. Setose bristle, from the middle of the superior series of the tenth foot of <i>Nereis patagonica</i> ,	× 400.
Fig. 2. A falcate bristle, from the inferior series of the same foot,	× 530.
Fig. 3. A setose bristle, from the middle of the superior series of the twelfth foot (in water) of <i>Nereis brasiliensis</i> ,	× 400.
Fig. 4. A falcate bristle, from the inferior border of the twelfth foot of the same,	× 530.
Fig. 5. Peculiar crenate dorsal bristle of <i>Staurocephalus atlanticus</i> ,	× 800.
Figs. 6, 7. Superior and inferior bristles of the ventral division of the foot of the foregoing,	× 450.
Fig. 8. One of the slender elongate bristles from the ventral division of the foot in the same,	× 350.
Fig. 9. One of the shorter dorsal bristles of <i>Staurocephalus australiensis</i> ,	× 250.
Fig. 10. Ventral bristle of the foregoing,	× 250.
Fig. 11. One of the superior winged bristles (in calcium chloride), from the tenth foot of <i>Lumbriconereis pettigrewi</i> ,	× 250.
Fig. 12. Bristle, from the middle of the same foot,	× 250.
Fig. 13. A posterior hook (in profile) of the same form, viewed laterally,	× 400.
Fig. 14. An antero-posterior view of a smaller hook from the same region,	× 400.
Fig. 15. Elongated hook, from the tenth foot of a variety of <i>Lumbriconereis pettigrewi</i> ,	× 250.
Fig. 16. Profile view of the upper hook, from a foot near the middle of the body of <i>Lumbriconereis bifurcata</i> ,	× 400.
Fig. 17. One of the third series (jointed hooks), from the anterior third of <i>Lumbriconereis japonica</i> ,	× 250.
Fig. 18. A posterior hook of <i>Lumbriconereis kerguelensis</i> ,	× 400.