

PLATE XIXA.

- Fig. 1. Brush-shaped bristle of *Nematonereis* (fragment) from Sea Point.  
     From the superior division of the foot, . . . . . × 350.
- Fig. 2. Articulated ventral bristle of the same species, . . . . . × 350.
- Fig. 3. Brush-shaped bristle, from the superior branch of *Nematonereis*,  
     from St. Vincent, . . . . . × 350.
- Fig. 4. Compound bristle, from the inferior division of the foot of the  
     same species, . . . . . × 350.
- Fig. 5. Long hook, from the inferior division of the foregoing, just after  
     development (*i.e.*, unworn), . . . . . × 350.
- Fig. 6. Dorsal bristle of *Eunice magellanica*, . . . . . × 210.
- Fig. 7. Brush-shaped bristle from the same group, . . . . . × 350.
- Fig. 8. Compound bristle, from the inferior division of the foot of the  
     foregoing, . . . . . × 210.
- Fig. 9. Strong bifid posterior hook from the same, one of the wings only  
     being indicated, . . . . . × 210.
- Fig. 10. Compound bristle, from the inferior division of the foot of *Eunice*  
     *prognatha*, . . . . . × 350.
- Fig. 11. Tip of stout blackish hook, from the posterior region of the same, . . . . . × 210.
- Fig. 12. Compound bristle, from the inferior group of the twentieth foot of  
     *Eunice torresiensis*, . . . . . × 350.
- Fig. 13. Posterior hook, from the same species, . . . . . × 210.
- Fig. 14. Compound bristle, from the twentieth foot of *Eunice ærstedii*, off  
     Marion Island, . . . . . × 350.
- Fig. 15. Compound bristle, from the same foot of a specimen of *Eunice*  
     *norvegica* from Bergen, Norway, . . . . . × 350.
- Fig. 16. Compound bristle, from the twentieth foot of *Eunice vittata*, . . . . . × 350.
- Fig. 17. Posterior hook, from the same species, . . . . . × 350.
- Fig. 18. Compound bristle, from the inferior border of the twentieth foot of  
     *Marphysa goodsiri*, . . . . . × 350.
- Fig. 19. Tip of the posterior hook of the same species, . . . . . × 350.
- Fig. 20. One of the lower (broader) bristles, from the dorsal group of the  
     foregoing, . . . . . × 350.
- Fig. 21. Superior bristle, from the twentieth foot of *Eunice cirrobranchiata*.  
     Not altogether a lateral view, . . . . . × 210.
- Fig. 22. Compound bristle from the same foot, slightly turned round, . . . . . × 350.
- Fig. 23. A more completely lateral view of a bristle from the same foot, the  
     shaft being somewhat injured by compression, . . . . . × 350.
- Fig. 24. Posterior hook of the same species, with the usual double wing, . . . . . × 210.