

PLATE XXA.

Fig. 1.	Dorsal bristle, from the anterior region of <i>Eunice kobeensis</i> ,	× 350.
Fig. 2.	Compound bristle, from the fortieth foot of the same, mounted in calcium chloride,	× 350.
Fig. 3.	Posterior hook of the foregoing,	× 350.
Fig. 4.	Dorsal bristle, from the tenth foot of <i>Eunice edwardsi</i> . Viewed nearly but not quite laterally,	× 350.
Fig. 5.	Brush-shaped bristle, from the same foot,	× 350.
Fig. 6.	Compound bristle of the same species, from the middle of the body,	× 350.
Fig. 7.	Inferior hook of the foregoing,	× 350.
Fig. 8.	Compound bristle, from the middle region of <i>Eunice aphroditois</i> ,	× 210.
Fig. 9.	Posterior hook, from the same form,	× 210.
Fig. 10.	Compound bristle, from the anterior third of <i>Eunice aphroditois</i> from Samboangan,	× 126.
Fig. 11.	Tip of a compound bristle, from the twentieth foot of <i>Eunice torquata</i> ,	× 350.
Fig. 12.	Fully developed and somewhat worn hook, from the posterior region of the same,	× 210.
Fig. 13.	Developing hook of the foregoing not yet projecting beyond the foot. The wing is imperfectly seen from the opacity of the surrounding tissue,	× 210.
Fig. 14.	Compound bristle, from one of the smaller examples of <i>Eunice elseyi</i> ,	× 350.
Fig. 15.	Similar bristle, from a large example of the foregoing species,	× 350.
Fig. 16.	Posterior hook from the latter specimen. The wing or guard, which is seldom complete, has been restored after an examination of several,	× 210.
Fig. 17.	Compound bristle, from the twentieth foot of <i>Nicidon balfouriana</i> ,	× 350.
Fig. 18.	Posterior hook, from the foregoing,	× 350.
Fig. 19.	Compound bristle, from the twentieth foot of <i>Eunice murrayi</i> ,	× 350.
Fig. 20.	Recently developed hook, from the posterior region of the same,	× 350.
Fig. 21.	Compound bristle (in water), from the twentieth foot of <i>Eunice mindanavensis</i> ,	× 350.
Fig. 22.	Inferior hook of the same,	× 210.
Fig. 23.	Compound bristle, the shaft being slightly turned round, from the twentieth foot of <i>Eunice equibranchiata</i> ,	× 350.
Fig. 24.	Inferior hook of the same,	× 210.
Fig. 25.	Bifid hook of <i>Macduffia bonhardi</i> ,	× 350.
Fig. 26.	Compound bristle of the foregoing,	× 350.