

PLATE XXIII.

- Fig. 1. Portion of a bristle, from the inferior tuft of the second foot of
Trophonia capensis, × 116.
- Fig. 2. Hook, from the ventral tuft of the tenth foot of the same, × 90.
- Fig. 3. Posterior bristle with hooked tip, from the foregoing, × 55.
- Fig. 4. Portion (near the tip) of a superior bristle, from the tenth foot of
Trophonia kerguelarum, × 350.
- Fig. 5. Hook, from the lower border of the tenth group of the same, × 90.
- Fig. 6. Upper hook, from the same series, × 90.
- Fig. 7. Portion of one of the larger frontal bristles of *Brada mammillata*,
with parasitic Infusoria, × 90.
- Fig. 8. Ventral hook, from the tenth foot of the same, × 90.
- Fig. 9. Basal region of one of the anterior bristles of *Brada whiteavesii*, × 350.
- Fig. 10. Anterior hook of the foregoing, × 90.
- Fig. 11. Portion of the basal region of a dorsal bristle, from the first foot
of *Trophonia wyvillei*. The proximal end is inferior, × 90.
- Fig. 12. Portion of the slender distal region of the same bristle where the
striæ are evident, × 90.
- Fig. 13. Bristle, from the ventral series of the tenth foot of the foregoing, × about 40.
- Fig. 14. Tip of one of the same series, × 90.
- Fig. 15. Extremity of a bristle, from the first series of *Buskiella abyssorum*, × 350.
- Fig. 16. Portion from the middle of one of the smaller forms from the
same group, × 350.
- Fig. 17. The same region of the bristle so as to show the length of the
divisions on each side of the transverse bar, × 55.
- Fig. 18. Tip of one of the ventral bristles from the foregoing, × 350.
- Fig. 19. Group of the fourth series of bristles in *Ranzania capensis*, × 350.
- Fig. 20. Bristle with expanded tip, from the same foot, × 350.