

PLATE XXIVA.

- Fig. 1. One of the first series of bristles in *Phyllochætopterus claparedii*,
in profile, × 350.
- Fig. 2. The same seen antero-posteriorly, × 350.
- Fig. 3. Strong brownish spine, from the middle of the third series of
bristles in the foregoing form, × 90.
- Fig. 4. Bristle, from the fifth series in this species, × 350.
- Fig. 5. Group of bristles, from the posterior region in the same example, × 350.
- Fig. 6. Hook, from about the fiftieth foot of *Scolecopsis cirrata*, var. 2,
from Sombrero and St. Thomas, × 350.
- Fig. 7. One of the larger hooks, from the tenth foot of *Prionospio capensis*,
seen partly but not quite in profile, × 350.
- Fig. 8. A smaller example from the same foot, exhibiting a full profile, × 350.
- Fig. 9. Fragment of a dorsal bristle, from the tenth series of *Cirratulus*
capensis, × 350.
- Fig. 10. Hook, from the posterior region of the same (in Farrant's solution), × 55.
- Fig. 11. Hook, from the posterior part of the fragment of *Cirratulus assimilis*, × 350.
- Fig. 12. Dorsal bristle, from the tenth foot of *Chætozone atlantica*, × 55.
- Fig. 13. Dorsal bristle, from the tenth foot of *Chætozone benthaliana*, × 55.
- Fig. 14. Posterior ventral bristles and spines in the same form, × 350.
- Fig. 15. Anterior hook of *Notomastus agassizii*, × 500.
- Fig. 16. Hook, from the second region of *Eunotomastus grubei*, × 500.
- Fig. 17. Hook of *Dasybranchus* sp. (fragment), × 500.
- Fig. 18. Dorsal bristle, from the eighth foot of *Nicomache capensis*, × 350.
- Fig. 19. Ventral hook, from the same foot of the foregoing, × 350.
- Fig. 20. Hook, from the eighth bristled foot of *Nicomache japonica*, × 350.
- Fig. 21. Posterior hook of *Nicomache* (?) *benthaliana*, × 350.