

1000 to 1200 fathoms.

Eumenia reticulata.
Melinna armandi.

Leæna neo-zealanica.
langerhansi.

1200 to 1500 fathoms.

Harmothoë benthaliana.
Polynoë synophthalma.
longipedata.
Nephtys phyllobranchia.
Laranda longa.
Lumbriconereis punctata.
ehlersi.
Eunice ærstedii (to 83 fathoms).
Aricia norvegica.
Chætozone benthaliana.
Notomastus agassizii.

Maldanella antarctica.
neo-zealanica.
Nicomache benthaliana.
Maldane sp.
Praxilla occidentalis.
Myriochele heeri.
Amphicteis gunneri.
Thelepus sp.
Terebellides stræmi.
Serpula philippensis.
Vermilia sp.

1500 to 2000 fathoms.

Chloenea atlantica.
Lætmonice producta, var. *wyvillei*
(and to 50 fathoms).
Lætmonice producta, var. *wille-*
moesii (to 700 fathoms).
Lagisca moseleyi.
crosetensis.
Dalhousia atlantica.
Nereis longisetis.
Nothria abranchiata.
armandi.

Nothria benthaliana.
Ephesia antarctica.
Trophonia wyvillei.
Maldane malmgreni.
Praxilla abyssorum.
Petta assimilis.
Amphicteis sarsi.
Grubianella antarctica, and var.
Melinnopsis atlantica.
Pista abyssicola.
Leæna antarctica.

2000 to 2500 fathoms.

Lumbriconereis abyssorum.
Nothria pycnbranchiata.
ehlersi.
Maldanella valparaisiensis.

Samythopsis grubei.
Eusamytha pacifica.
Melinna pacifica.
Euthelepus chilensis.

Placostegus mörchii.