

GEOGRAPHICAL DISTRIBUTION.

The following table exhibits the geographical distribution and the occurrence of the species described. I have also noted the localities where those species already known have been previously found.

Species observed in the Challenger and "Porcupine" Expeditions.						Species previously observed.	
Name.	Station.	"Porcupine" Voyage.	Locality.	Depth in Fathoms.	Character of the Bottom.	Depth.	Locality.
<i>Sternaspis princeps</i> , n. sp.	167	...	lat. 37° 34' S. long. 179° 22' E.	700	Blue mud.
<i>Echiurus uncinatus</i> , v. Drasche.	Japan.	Japan.
<i>Thalassema faxa</i> , n. sp.	...	P.	lat. 60° 34' N. long. 4° 40' E.	560
<i>Thalassema baronii</i> , Groeff.	Bahia.	7-20
<i>Bonellia suhmii</i> , n. sp.	47	...	Nova Scotia, lat. 41° 15' N. long. 65° 45' W.	1340	Blue mud.
<i>Phascolosoma vulgare</i> , Blainville.	30	P.	lat. 36° 15' N. long. 6° 32' W.	386	...	Near low water mark.	West coast of Europe, Mediterranean, Red Sea.
<i>Phascolosoma vulgare</i> , var. <i>astuta</i> , nov.	90	P.	lat. 59° 41' N. long. 7° 34' W.	458
<i>Phascolosoma pudicum</i> , n. sp.	Kerguelen.	10-120
<i>Phascolosoma capense</i> , Teuscher.	Sea Point, near Cape Town.	Shallow water.	Cape.
<i>Phascolosoma prioki</i> , Sluiter.	214	...	lat. 4° 33' N. long. 127° 6' E.	500	Blue mud.	...	Batavia.
<i>Phascolosoma catharinae</i> , Fritz Müller.	323	...	lat. 35° 39' S. long. 50° 47' W.	1900	Desterro.
<i>Phascolosoma flagriferum</i> , n. sp.	241	...	lat. 35° 41' N. long. 137° 42' E.	2800
	45	...	lat. 37° 25' N. long. 71° 43' W.	1700
<i>Dendrostoma blandum</i> , Selenka and de Man.	Yokohama.	5-25	Japan.
<i>Phascolion lutense</i> , n. sp.	lat. 62° 26' S. long. 95° 44' E.	1975	Diatom ooze.
			lat. 53° 55' S. long. 108° 35' E.	1950	Diatom ooze.
<i>Phascolion tubicola</i> , Verrill.	49	...	lat. 4° 21' S. long. 129° 7' E.	85	Gravel, stones.	Frequently in deep water.	New England.
<i>Phascolion botulus</i> , n. sp.	195	...	lat. 4° 21' S. long. 129° 7' E.	1425	Blue mud.
<i>Phascolion strombi</i> (Montagu).	...	P.	Killibegs Harbour, Ireland.	North Atlantic.
<i>Phascolion squamatum</i> , Koren and Danielssen.	6 10 36 47	P.	...	358-725	...	100-200	North Sea, Lofoten, &c.
<i>Aspidosiphon speculator</i> , n. sp.	St. Vincent (Cape Verdes).	Shallow water.
<i>Aspidosiphon truncatus</i> , Keferstein.	201	...	lat. 7° 3' N. long. 121° 48' E.	82	Stones, gravel.	...	Panama, Mauritius.
<i>Phymosoma japonicum</i> , Grube.	Port Jackson.	Shallow water.	Japan.
<i>Phymosoma scolops</i> , Selenka and de Man.	208	...	lat. 11° 37' N. long. 123° 32' E.	18	Blue mud.	...	Philippines, Singapore, Mozambique, Red Sea.
<i>Sipunculus nudus</i> , Linné.	30	P.	lat. 36° 18' N. long. 6° 52' W.	386	Mediterranean, West coast of Europe, West Indies, Philippines.
	22	...	lat. 56° 8' N. long. 13° 34' W.	1263
<i>Chaetoderma militare</i> , n. sp.	210	...	lat. 9° 28' N. long. 123° 45' E.	375	Blue mud.