

| | PAGE | | PAGE |
|--|----------|---|------------|
| <i>monacaria</i> , Jæger (Holothuria), . | 172, 217 | <i>okeni</i> , Bell (Thyone), | 139 |
| <i>monacaria</i> , Lesson (Psolus), . | 217 | <i>Oligotrochus</i> , Sars, | 37 |
| Monooyolia , Lampert (Group), . | 266 | <i>Oncinolabes</i> , Brandt, | 19, 20, 22 |
| <i>monosticha</i> , Haacke (Holothuria), . | 239 | <i>oolitica</i> , Semper (Haplodactyla), . . | 51, 53 |
| <i>monotuberculata</i> , Quoy and Gaimard (Holothuria), . . | 197 | <i>oolitica</i> , Selenka (Molpadia), . . | 51, 53 |
| <i>moseleyi</i> , n. sp. (Stichopus) . | 165 | <i>ooliticum</i> , Pourtalès (Chirodota), . . | 53 |
| <i>montagui</i> , Bell (Cucumaria), . | 116 | <i>ooliticum</i> , Danielssen and Koren (Trochostoma), | 53 |
| <i>montagui</i> , Fleming (Holothuria), . | 106 | <i>ooplax</i> , von Marenzeller (Synapta), . | 25 |
| <i>Mülleria</i> , Jæger, . . | 134, 197 | <i>operculatus</i> , Pourtalès (Cuvieria), . | 130 |
| <i>multipes</i> , n. sp. (Cucumaria), . | 72 | <i>operculatus</i> , Verrill (Lepidopsolus), . | 130 |
| <i>muricata</i> , Studer (Trachythylene), . | 139 | <i>operculatus</i> , Pourtalès (Psolus), . . | 87 |
| <i>murrayi</i> , n. sp. (Holothuria), . | 185, 187 | <i>ophidiana</i> , Quoy and Gaimard (Holothuria), . | 240 |
| <i>murrayi</i> , n. sp. (Psolus), . . | 85 | <i>Orcula</i> , Troschel, | 97, 149 |
| <i>murrayi</i> , Bell (Stereoderma), . | 142 | <i>ornatus</i> , Verrill (Lissothuria), . . | 127 |
| <i>musculosa</i> , Semper (Thyone), . | 141 | <i>ovata</i> , Risso (Holothuria), . . | 240 |
| <i>musculosa</i> , Ayres (Thyonidium), . | 141 | <i>ovulum</i> , Selenka (Stolus), . . | 138 |
| <i>musculus</i> , Semper (Haplodactyla), . | 50 | <i>ovulum</i> , Semper (Thyone), . . | 138 |
| <i>musculus</i> , Risso (Molpadia), . | 50 | <i>Palopatides</i> , n. gen., | 154 |
| <i>Myriotrochus</i> , Steenstrup, . . | 37 | <i>pallidus</i> , Ayres (Trochinus), . . | 34 |
| <i>naso</i> , Haacke (Stichopus), . | 191 | <i>panaensis</i> , Semper (Chirodota), . . | 37 |
| <i>naso</i> , Semper (Stichopus), . | 192 | <i>panamensis</i> , Verrill (Pentacta), . . | 116 |
| <i>natans</i> , Sars (Holothuria, Stichopus), . | 193 | <i>panamensis</i> , Théel (Cucumaria), . . | 116 |
| <i>neillii</i> , Leslie and Herdman (Cucumaria), . | 116 | <i>pandanensis</i> (= boholensis, var.), Semper (Psolus), | 128 |
| <i>neillii</i> , Fleming (Holothuria), . | 116 | <i>papillosa</i> , O. F. Müller (Holothuria), . . | 134 |
| <i>neglectum</i> , Haacke (Labidodemas), . | 218 | <i>papillosa</i> , Johnston (Mülleria), . . | 134 |
| <i>niger</i> , Kinahan (Cucumaria), . | 238 | <i>papillosa</i> , Forbes (Thyone), . . | 134 |
| <i>nigra</i> , Peach (Holothuria), . | 238 | <i>papuensis</i> (= fusus, var.), nov. (Thyone), . | 92 |
| <i>nigra</i> , Semper (Synapta), . | 20 | <i>paradoxa</i> , Selenka (Holothuria), . . | 206 |
| <i>nigricans</i> , Selenka (Cucumaria), . | 104 | <i>paradoxus</i> , Lampert (Stichopus), . . | 265 |
| <i>nigricans</i> , Brandt (Cladodactyla, subg. Polyclados), . | 104 | <i>pardalis</i> , Selenka (Holothuria), . . | 224 |
| <i>nigricans</i> , Stimpson (Pentacta), . | 104 | <i>parva</i> , Krauss, Lampert (Holothuria), . . | 264 |
| <i>nobilis</i> , Ludwig (Cucumaria), . | 114 | <i>parva</i> , Ludwig (Cucumaria), . . | 111 |
| <i>nobilis</i> , Lampert (Semperia) = <i>Cucumaria nobilis</i> . | | <i>parva</i> , Lampert (Semperia) = <i>Cucumaria parva</i> . | |
| <i>nobilis</i> , Selenka (Mülleria), . | 198 | <i>parva</i> (= murrayi, var.), nov. (Holothuria), . | 186 |
| <i>notabilis</i> , Ludwig (Holothuria), . | 228 | <i>parvula</i> , Verrill (Actinopyga), . . | 199 |
| <i>obesa</i> , Selenka (Mülleria), . | 200 | <i>parvula</i> , Haacke (Mülleria), . . | 219 |
| <i>obscura</i> , Verrill (Actinopyga), . | 238 | <i>parvula</i> , Selenka (Mülleria), . . | 199 |
| <i>obscura</i> , Lesueur (Holothuria), . | 238 | <i>parvum</i> , Ludwig (Thyonidium), . . | 147 |
| <i>obunca</i> , Lampert (Cucumaria), . | 265 | <i>Pattalus</i> , Selenka, | 144 |
| <i>occidentale</i> , Ludwig (Thyonidium), . | 147 | <i>pauper</i> , Selenka (Embolus), . . | 56 |
| <i>occidentalis</i> , Ludwig (Holothuria), . | 228 | <i>PEDATA</i> , Claus (Order), | 57 |
| <i>oceania</i> , Lesson (Holothuria), . | 27 | <i>pedata</i> , Semper (Thyone), . . | 140 |
| <i>oceania</i> , Jæger (Synapta), . | 27 | <i>pellucida</i> , Blainville (Cucumaria), . . | 34 |
| <i>ocellata</i> , Jæger (Bohadschia), . | 239 | <i>pellucida</i> , Brandt (Dactylota), . . | 34 |
| <i>ocellata</i> , Semper (Holothuria), . | 178, 239 | <i>pellucida</i> (= molpadioides, var.), Semper (Haplodactyla), . . | 50 |
| <i>Ocnus</i> , Forbes, . . | 75, 117 | | |