

PLATE IX.

- Fig. 1. *Stichopus möbii*, Semper. *a* and *b*, under view and side view of the common tables ; *c*, end of the spire of the same tables, seen from above ; *d*, side views of larger tables ; *e*, under views of larger tables ; *f*, supporting rods from the pedicels ; *g*, C-shaped body.
- Fig. 2. *Holothuria murrayi*, var. *parva*, nov. *a*, disk of the common form of tables, seen from the under surface ; *b*, disk of the rare tables, seen from the under surface ; *c*, side view of a table ; *d*, end of the spire of the tables.
- Fig. 3. *Holothuria murrayi*, n. sp. (var.?). Calcareous ring, three times the natural size.
- Fig. 4. *Thyonidium cebuense*, Semper. *a*, calcareous ring, about seven times the natural size ; *b*, *c*, *d*, diverse views of small tables ; *e*, a large table seen from above ; *f*, rod-like tables from the pedicels.
- Fig. 5. *Cucumaria mirabilis*, n. sp. *a*, calcareous ring, about twenty times the natural size ; *b*, *c*, *d*, diverse views of tables ; *e*, *f*, diverse views of supporting rods from the ambulacral appendages.
- Fig. 6. *Psolus disciformis*, n. sp. *a*, *b*, plates from the sole ; *c*, rods from the pedicels.
- Fig. 7. *Pælopatides confundens*, n. sp. *a*, dorsal view ; *b*, ventral view, natural size.