

Lamellæ mandibulares validiores, structura uniformi (quasi homogena). Dens medianus radulæ basi (fere semper) bifidus; dens lateralis magnus, dorso fissus, hamo curvato marginibus serrulato. Proventriculus non deest. Pars inferior varis deferentis in cavitate abdominali pro parte libera.

Testa interna calcarea, lactea, nonnihil diaphana, lævis, obovata, spirata, anfractibus vix tribus; spira ad $\frac{1}{4}$ longitudinis sita, prominenti, acutiuscula, anfractibus vix duobus; apertura ampla, ovata, patente usque in fundum spiræ, columella antice sinu levi excisa.

- | | |
|---|---|
| <p>1. <i>Marsenia tentaculata</i> (Mtg.).
Mare atlant. orient.</p> <p>2. <i>Marsenia producta</i>, Leach.
Mare atlant. orient.</p> <p>3. <i>Marsenia perspicua</i>¹ (Linné).
Var. (?) Morelli, Delle Chiaje.
Mare mediterr.</p> <p>4. <i>Marsenia dubia</i>, n. sp.
Mare atlant. orient. (Cape Verde).</p> <p>5. <i>Marsenia pellucida</i> (Verrill).
Mare atlant. occid.</p> <p>6. <i>Marsenia stearnsii</i> (Dall).
Mare pacific. (Californ.).</p> <p>7. <i>Marsenia orbiculata</i> (Dall).
Mare pacific. (Californ.).</p> <p>8. <i>Marsenia latens</i> (O. Fr. Müller).
Mare atlant. or. sept.</p> <p>9. (?) <i>Marsenia rangii</i>, Bgh.
Mare antillense.</p> <p>10. <i>Marsenia minuta</i> (H. Adams).
Mare rubrum.</p> | <p>11. <i>Marsenia cabulana</i>, Bgh.; n. sp.
Mare philippin.</p> <p>12. <i>Marsenia affinis</i>, Bgh., n. sp.
Mare philippin.</p> <p>13. <i>Marsenia gemma</i>, Bgh., n. sp.
Mare philippin.</p> <p>14. <i>Marsenia indecora</i>, Bgh., n. sp.
Mare philippin.</p> <p>15. <i>Marsenia isabellina</i>, Bgh., n. sp.
Mare philippin.</p> <p>16. <i>Marsenia dlegænsis</i> (Dall), n. sp.
Mare pacific.</p> <p>17. <i>Marsenia ophione</i> (Gray).
Mare pacific. (Nov. Zel.).</p> <p>18. <i>Marsenia Kerguelenensis</i> (Studer).
Mare pacific. australe.</p> <p>19. <i>Marsenia</i> (?) <i>translucida</i>, Blv.
<i>Hab. ?</i></p> <p>20. <i>Marsenia depressa</i>, Wood.
Crag (Sutton).</p> |
|---|---|

II. *Chelyonotus* (Swains.), Bergh.

Animalia diœcia, Marseniis propriis similia; sed plica respiratoria producta, apice interdum bifida. Lamellæ branchiales fere ut in Marseniis.—Lamellæ mandibulares et armatura lingualis fere ut in Marseniis, sicut proventriculus. Vas deferens in cavitate abdominali non liberum sed in pariete abdominali omnino inclusum.

Testa interna fere omnino ut in Marseniis sed magis calcarea.

- | | |
|---|--|
| <p>1. <i>Chelyonotus semperi</i>, Bgh.
Mare philippin.</p> <p>2. <i>Chelyonotus tonganus</i> (Quoy et Gaim.).
Mare indicum.</p> | <p>3. <i>Chelyonotus tuberosus</i> (Stimpson).
Mare chinense.</p> <p>4. <i>Chelyonotus</i> (?) <i>punctatus</i> (Stimpson).
Mare chinense.</p> |
|---|--|

¹ The species indicated by larger italic type have been anatomically examined by the author. The bracket indicates that the species enclosed by it are nearly related to each other.