7. Phos smithi, n. sp. (Plate XVII. fig. 7).

Station 122. September 10, 1874. Lat. 9° 5' S., long. 34° 50' W. Off Pernambuco. 350 fathoms. Mud.

Shell.—Long, narrow, fusiform, closely and narrowly ribbed, spiralled, with a high narrow spire of flatly convex whorls, a small sharp apex, a shortish conical base, terminating in a short, lop-sided, straight snout. Sculpture: Longitudinals—there are on the last whorl 18 or 19 ribs, which diminish slowly in number up the spire; they stretch from suture to suture, are more or less continuous from whorl to whorl, and extend to the snout; they are narrow but well rounded, though on the last whorl they tend a little to spread out and flatten; the furrows between them are rather broader than they, especially on the body-whorl, for towards the apex they are crowded; on each whorl one or two of these ribs swell into a slightish varix: the entire surface is densely covered with fine hair-like lines of growth. Spirals—the whole shell is scored with rounded threads a good deal finer than the ribs; of these there are 2 on the first regular whorl, 7 to 8 on the penultimate, and about 16 on the body-whorl; in crossing the ribs these threads rise into broadish compressed white tubercles; the furrows between the threads are more than double the breadth of these, but generally have a finer thread in the bottom; two of the threads at the periphery, and the two on the snout, are stronger than the rest; the point of the snout is scored with the old canal scar, and is streaked obliquely with narrowish threads. Colour an ochrey white, stained with three bands of pale chestnut brown. Spire high, narrow, conical. Apex is sharply conical, and consists of 4 small white carinated angulated whorls (the extreme tip is broken, but was evidently minute), of which the last is feebly ribbed longitudinally. Whorls 8 below those of the apex; they are short and convex, and of very slow increase; the last is small and narrow, and has a short rounded base, ending in a smallish, lop-sided, slightly twisted, barely reverted, emarginate snout. Suture a little impressed, rather oblique. Mouth small, oval, pear-shaped, prolonged into a very short oblique canal. Outer lip thin, flatly convex in its curve, patulous in front, where it becomes concave in its curve, and then it runs oblique and direct down the short canal. Inner lip narrowly excavated into the texture of the shell; it is flat on the body, concave at the base of the pillar, which is short, straight and feebly double-toothed at its point, where it is obliquely cut off and twisted. H. 1.25 in. B. 0.45. Penultimate whorl, height 0.23. Mouth, height 0.52, breadth 0.25.

This species a good deal resembles *Phos cumingii*, Reeve, in form, but the sculpture is quite different. I have given it the name of Mr Edgar A. Smith, whose ever kind help I have repeatedly had to appeal to.