

18. *Rissoa (Alvania) trajectus*, n. sp.

19. *Rissoa (Alvania) lusciniæ*, n. sp.

C. Spiralled strongly (*Onoba*), i.e., in which the spiral threads are the striking feature.

20. *Rissoa (Onoba) fenestrata*, Krauss.

21. *Rissoa (Onoba) arenaria*, Mighels.

22. *Rissoa (Onoba) striata*, (J. Ad).

23. *Rissoa (Onoba) Watsoni*, Schwartz.

24. *Rissoa (Onoba) brachia*, n. sp.

25. *Rissoa (Onoba) ædonis*, n. sp.

26. *Rissoa (Onoba) mercurialis*, n. sp.

D. Spiralled weakly (*Ceratia*), i.e., in which there are spiral threads, but these not strong.

27. *Rissoa (Ceratia) glaphyra*, n. sp.

28. *Rissoa (Ceratia) pachia*, n. sp.

29. *Rissoa (Ceratia) transenna*, n. sp.

30. *Rissoa (Ceratia) macra*, n. sp.

E. Whorls flat and smooth (*Cingula*).

31. *Rissoa (Cingula) glabrata*, v. Mühlf.

32. *Rissoa (Cingula) rustica*, n. sp.

33. *Rissoa (Cingula) scopulorum*, n. sp.

34. *Rissoa (Cingula) alvearium*, n. sp.

F. Whorls rounded and smooth (*Setia*).

35. *Rissoa (Setia) albugo*, Wats.

36. *Rissoa (Setia) callosa*, Manzoni.

37. *Rissoa (Setia) tenuisculpta*, Wats.

38. *Rissoa (Setia) marionensis*, n. sp.

39. *Rissoa (Setia) principis*, n. sp.

40. *Rissoa (Setia) australis*, n. sp.

41. *Rissoa (Setia) philomelæ*, n. sp.

42. *Rissoa (Setia) quisquiliarum*, n. sp.

43. *Rissoa (Setia) edwardiensis*, n. sp.

44. *Rissoa (Setia) sinapi*, n. sp.

45. *Rissoa (Setia) triangularis*, n. sp.

G. Mouth separated from body-whorl (*Scrobs*).

46. *Rissoa (Scrobs) scrobiculator*, n. sp.

47. *Rissoa (Scrobs) badia*, n. sp.

A. Ribbed (*Rissoa*), i.e., of which the longitudinal ribs are the striking feature.

1. *Rissoa costulata*, Alder.

Rissoa costulata, Alder, Ann. and Mag. Nat. Hist., 1844 (ser. 1), vol. xiii. p. 324, pl. viii. figs. 8, 9.

„ „ Forbes and Hanley, Brit. Moll., vol. iii. p. 103, pl. lxxvii. figs. 4, 5.

„ *subcostulata*, Schwartz v. Mohrenstern, *Rissoa*, p. 41, sp. 32, pl. iii. fig. 32.

„ *costulata*, Jeffreys, Brit. Conch., vol. iv. p. 35, vol. v. p. 208, pl. lxxviii. fig. 1.

„ „ Weinkauff, Conch. Mittelmeer., vol. ii. p. 297, sp. 20.

„ „ Watson, Proc. Zool. Soc. Lond., 1873, p. 378, pl. xxxv. fig. 15.

„ „ Reeve, Conch. Icon., vol. xx. pl. iii. fig. 29.

„ „ Monterosato, Enumerazione, p. 24.

„ „ Seguenza, Form. Terz. Calabria, pp. 268, 355.

„ „ Weinkauff, Conch. Cab. (ed. Küster), p. 139, sp. 37, pl. x. figs. 1-3.

Apicularia guerini, Monterosato, Nomenclatura, p. 56.

February 10, 1873. Tenerife, Canaries. 70 fathoms.

Habitat.—Great Britain to the Mediterranean (Jeffreys); Madeira (Watson).

Fossil.—From the Middle Pliocene onwards (Seguenza).

The Marquis de Monterosato holds this as *Rissoa guerini*, Recluz, which the Baron Schwartz von Mohrenstern considers a distinct species. As I have remarked elsewhere and below in connection with *Rissoa parva*, Scacchi, the whole group absolutely requires revision, and nothing but worse confusion can result from mere partial meddling with it in the meantime. The *Rissoa costulata* of Risso (1826) is a synonym of the earlier *Rissoa variabilis*, v. Mühlf. (1818), and is therefore out of reckoning.