

STATION 233.—Bay of Kobé, Japan, May 17–19, 1875; lat. $34^{\circ} 39'$ N., long. $135^{\circ} 14'$ E.; 8 fathoms; mud; surface temperature, $62^{\circ} 3$. Dredge.

<i>Inioteuthis morsei.</i>		<i>Sepia kobiensis</i> , n. sp.
		<i>Loligo kobiensis</i> , n. sp.

STATION 233c.—Inland Sea, Japan, May 28, 1875; lat. $34^{\circ} 18'$ N., long. $133^{\circ} 21'$ E.; 12 fathoms; blue mud; bottom temperature, $59^{\circ} 9$; surface temperature, $66^{\circ} 8$. Trawl.

<i>Sepiella maindroni</i> (?).		<i>Loligo kobiensis</i> , n. sp.
		<i>Todarodes pacificus.</i>

STATION 237.—South-east of Nosima, Japan, June 17, 1875; lat. $34^{\circ} 37'$ N., long. $140^{\circ} 32'$ E.; 1875 fathoms; blue mud; bottom temperature, $35^{\circ} 3$; surface temperature, 73° . Trawl.

<i>Octopus januarii</i> , n. sp.		<i>Promachoteuthis megaptera</i> , n. sp.
----------------------------------	--	---

Reefs, Honolulu, Sandwich Islands.

<i>Octopus marmoratus</i> , n. sp.

Pacific Ocean, south of the Sandwich Islands, September 2, 1875; lat. $5^{\circ} 54'$ N., long. $147^{\circ} 2'$ W. (Taken from the stomach of a shark.)

<i>Chiroteuthis</i> (?) sp.

STATION 271.—Central Pacific, September 6, 1875; lat. $0^{\circ} 33'$ S., long. $151^{\circ} 34'$ W.; 2425 fathoms; Globigerina ooze; bottom temperature, 35° ; surface temperature, $78^{\circ} 7$. Trawl.

<i>Enoplateuthis margaritifera.</i>

South Pacific, between the Sandwich Islands and Tahiti, September 15, 1875; lat. $12^{\circ} 8'$ S., long. $150^{\circ} 13'$ W.; surface temperature, 80° .

<i>Tremoctopus quoyanus.</i>

Off Valparaiso, October 21, 1875; surface to 30 fathoms.

<i>Onychoteuthis</i> sp.

South Pacific, November 1, 5, 7, and 11, 1875; about lat. $38^{\circ} 7'$ S., long. $94^{\circ} 4'$ W.; surface.

<i>Tracheloteuthis</i> sp.
