

Dosidicus eschrichtii.
Tracheloteuthis riisei (also A. XII.).
Abralia polyonyx.
 „ *veranyi.*
 „ *oweni.*
Enoplateuthis margaritifera (also C.).
Verania sicula.

Ancistroteuthis lichtensteinii.
Teleoteuthis krohnii.
Gonatus fabricii, (also A. C. II. XVII.).
Chiroteuthis veranyi.
Histioteuthis bonelliana.
 „ *rupPELLI.*
Doratopsis vermicularis.

VI. THE WEST AFRICAN REGION.

The district thus named extends from the Canaries to about the Tropic of Capricorn.

Octopus vulgaris (also I. III. IV. V. VII.
 VIII. IX. X. XIII.).
 „ *tuberculatus* (also III. V.).
 † „ *occidentalis* (also III.).
 † „ *granulatus* (also III. IV. VII.
 IX. XV.).

Octopus macropus (also V. VIII. IX. X.).
Sepiola rondeleti (also I. IV. V.).
Sepia officinalis (also I. IV. V.).
 „ *hierredda.*
 „ *bertheloti.*
Sepiella ornata.

VII. THE SOUTH AFRICAN REGION.

A considerably greater area has been allotted to this province than is given to the corresponding one in Dr. Fischer's arrangement. It has been allowed to extend from the Cape as far as the Red Sea and to include the Madagascar and Mauritius, as well as the islands of the South Atlantic and Southern Oceans, the Tristan and Prince Edward groups, with the Kerguelen and Heard Islands.

† *Argonauta argo* (also A. C. II. IV. V. IX.
 XI.).
 „ *tuberculata* (also C. IX. XI.
 XII.).
Octopus vulgaris (also I. III. IV. V. VI.
 VIII. IX. X. XIII.).
 † „ *verrucosus.*
 † „ *granulatus* (also III. IV. VI. IX.
 XV.).
 ? „ *horridus* (also VIII.).
 „ *aranea* (also XIII.).
 „ *levis.*
Cistopus indicus (also IX.).
Sepiola stenodactyla.

Sepia vermiculata.
 „ *zanzibarica.*
 „ *venusta.*
 „ *capensis.*
 „ *tuberculata.*
Hemisepiopus typicus.
Sepioteuthis mauritiana.
 „ *loliginiformis* (also VIII.).
 „ *madagascariensis.*
Loligo reynaudii.
Monchezia sancti-pauli.
Enoplateuthis hoylei.
Pyrgopsis rhynchophorus.
Taonius (?) maximus.