

<i>Sepioteuthis lunulata</i> (also XI.).		<i>Tracheloteuthis riisei</i> (also A. V.).
† „ <i>lessoniana</i> (also IX. X. XII.).		<i>Nautilus pompilius</i> (also XI.).
<i>Loligo pfefferi</i> .		„ <i>scrobiculatus</i> .
„ <i>spectrum</i> .		„ <i>macromphalus</i> .
		„ <i>umbilicatus</i> .

XIV. THE CALIFORNIAN REGION.

This region I regard as stretching from the peninsula of Alaska to the Isthmus of Panama; probably it will eventually be necessary to subdivide it, but so few forms have been described from that coast, that this course hardly seems advisable at present. From the coast between Alaska and Kamtschatka no Cephalopods are known to me; probably they will be found like the other Mollusca from that region to be of Arctic types.

<i>Argonauta pacifica</i> .		<i>Octopus punctatus</i> (also IX. X.).
„ <i>expansa</i> .		<i>Loliolus steenstrupi</i> .
<i>Octopus bimaculatus</i> .		<i>Onychoteuthis lobipennis</i> .
		<i>Ancistroteuthis robusta</i> .

XV. THE PERUVIAN REGION.

The northern boundary of this province may be taken at the Isthmus of Panama and the southern at about the northern limit of Patagonia.

<i>Octopus granulatus</i> (also III. IV. VI. VII. IX.).		<i>Loligo gahi</i> (also III. XVI.).
„ <i>fontanianus</i> (also XVI.).		<i>Steenstrupiola chilensis</i> . ¹
		<i>Onychoteuthis brachyptera</i> . ¹
		<i>Taonius schneehageni</i> . ¹

XVI. THE PATAGONIAN REGION.

This region includes the extremity of South America, both on the eastern and western coasts.

<i>Octopus fontanianus</i> (also XV.).		<i>Rossia sublevis</i> (also II.).
„ <i>tehuelchus</i> (also III.).		<i>Loligo gahi</i> (also III. XV.).
† <i>Rossia patagonica</i> .		„ <i>patagonica</i> .
		<i>Onychoteuthis ingens</i> .

¹ Probably these species are Oceanic.