

TABLE VII.—Species recorded from between 2000 and 3000 fathoms.

Species.	Challenger.	Other British Expeditions.	American Expeditions.	Other Sources and Remarks.	Also in Table
<i>Cirroteuthis magna</i> , n. sp., . . .	2225				VI.
„ <i>pacifica</i> , n. sp., . . .	2440				
„ <i>megaptera</i> , . . .			2512, 2574		VI. VI.
<i>Eledone rotunda</i> , n. sp., . . .	2225				
<i>Eledonella pygmaea</i> , . . .			[2949]		
<i>Mastigoteuthis agassizii</i> , . . .			2515		V. VI. VI.
<i>Enoploteuthis margaritifera</i> , . . .	[2425]			Very doubtful.	VI. IV. VI.
<i>Chiroteuthis lacertosa</i> , . . .			2098, 2949		IV. VI.
<i>Calliteuthis reversa</i> , . . .			2369		IV. V. VI.
<i>Histiopsis atlantica</i> , n. sp., . . .	2025			Possibly surface.	
<i>Taonius suhmi</i> , n. sp., . . .	[2150]			Surface.	I.

Some of the genera enumerated in the above list merit a brief discussion, and among these perhaps the most interesting is *Cirroteuthis*, whose remarkable history has already been alluded to (p. 55); the type species, described in 1836, was the only one known till 1883, since which time six species have been brought to our knowledge in addition to the two closely allied if not identical genera, *Stauroteuthis* and *Opisthoteuthis*. The following table gives all the particulars which have been published regarding the habitat of each of these forms, as well as the localities of some specimens too fragmentary for identification.

Species.	Locality.	Depth in Fathoms.	Temperature.		Remarks.
			Bottom ° F.	Surface ° F.	
<i>Cirroteuthis mulleri</i> , Eschricht,	Jakobshavn, Greenland.	...	...	...	No information as to depth.
„ <i>umbellata</i> , Fischer, }	Off Azores and West Coast of Africa.	616 to 1206 }	...	...	
„ <i>magna</i> , Hoyle, . . .	Southern Ocean.	1375	35·6	43·0	
„ „ . . .	Off Valparaiso.	2225	35·6	59·0	
„ <i>pacifica</i> , Hoyle, . . .	Between Papua and Australia.	2440	35·8	80·0	Station Book says "surface."
„ <i>meangensis</i> , Hoyle, }	Off Meangis Islands, North of New Guinea.	500 }	41·8	80·5	
„ „ . . . }	Off Kermadec Islands, South Pacific.	600 }	39·5	66·5	
„ sp.?, . . . . .	20° West of Patagonia.	...	...	...	Floating on surface, dead and mutilated.
„ <i>plena</i> , Verrill, . . .	5° East of Chesapeake Bay.	1073	38·0	73·0	
„ <i>megaptera</i> , Verrill, . . .	A little further east.	2512	37·0	78·0	Two specimens.
„ „ „ . . . . .	A little further east still.	2574	37·0	79·0	„ „
„ sp.?, . . . . .	A little further north.	1054	38·0	74·0	
„ eggs of, . . . . .	More than six localities in the same region.	428 to 1106	38·0 to 40·0	71·0 to 74·0	
<i>Stauroteuthis syrtensis</i> , Verrill, }	30 miles east of Sable Island, Nova Scotia.	250 }	...	...	
„ „ . . . . .	Off Martha's Vineyard.	523	39·0	68·0	
„ „ . . . . .	Near same locality.	1346	...	...	
<i>Opisthoteuthis agassizii</i> , Verrill, }	Off Grenada.	291 }	...	...	
„ „ . . . . .	Off Martha's Vineyard.	1058	38·0	74·0	