PLATE XVIII.

		Figs. 1-6. Sepia esculenta, n. sp.,	PAGE 129
Fig.	1.	Ventral aspect of the shell of the female; natural size.	
Fig.		Dorsal aspect of the shell of the female; natural size.	
Fig.	3.	Ventral aspect of the shell of the male; natural size.	
Fig.	4.	Part of the margin of a sucker from one of the sessile arms. The upper part of the drawing shows the smooth horny ring, while below it is the papillary area; highly magnified.	
Fig.	5.	Sucker from one of the sessile arms; magnified about 8 diameters.	
Fig.	6.	The parts surrounding the mouth of the male, showing the bases of the arms, the hectocotylisation of the left ventral one, the buccal membrane, and the inner and outer lips; natural size.	J¥
		Figs. 7-14. Sepia kobiensis, n. sp.,	142
Fig.	7.	Dorsal view of the specimen; natural size.	
Fig.	8.	Ventral view of the specimen; natural size. Showing the fins terminating on the ventral surface some distance from the extremity of the animal.	٠
Fig.	9.	Club of the left tentacle; magnified 7 diameters.	
Fig.	10.	A sucker from one of the sessile arms; magnified 20 diameters.	,
Fig.	11.	Another view of a similar sucker, in which the distal depression is not marked; magnified 20 diameters.	
Fig.	12.	Lateral view of the shell; natural size.	
Fig.	13.	Dorsal view of the shell; natural size.	

Fig. 14. Ventral view of the shell; natural size.