

PLATE XIX.

Figs. 1-13. *SEPIA SULCATA*, n. sp., . . .

PAGE
137

- Fig. 1. Ventral aspect of the specimen ; natural size.
- Fig. 2. Dorsal aspect of the specimen ; natural size.
- Fig. 3. The left tentacular club ; magnified 7 diameters.
- Fig. 4. The hectocotylised arm ; magnified 2 diameters.
- Fig. 5. A transverse section of the same, to show the groove and the rounded fillet lying in it ; magnified 3 diameters.
- Fig. 6. A tentacular sucker ; magnified 40 diameters.
- Fig. 7. A sucker from one of the sessile arms ; magnified 40 diameters.
- Fig. 8. A portion of the papillary area ; magnified 200 diameters.
- Fig. 9. The ventral mandible ; natural size.
- Fig. 10. The dorsal mandible ; natural size.
- Fig. 11. Ventral aspect of the shell ; natural size.
- Fig. 12. Lateral aspect of the shell ; natural size.
- Fig. 13. Dorsal aspect of the shell ; natural size.

Figs. 14-24. *SEPIA ELLIPTICA*, n. sp., . . .

131

- Fig. 14. Dorsal view of a specimen ; natural size.
- Fig. 15. Ventral view of a specimen ; natural size.
- Fig. 16. Inner aspect of the hectocotylised arm ; magnified 2 diameters.
- Fig. 17. A sucker from a sessile arm, with smooth horny ring ; magnified 20 diameters.
- Fig. 18. A portion of the papillary area, the upper margin is the horny ring ; magnified 200 diameters.
- Fig. 19. A somewhat compressed sucker from a sessile arm, showing slight tothing of the distal margin of the horny ring ; magnified 20 diameters.
- Fig. 20. Ventral mandible ; natural size.
- Fig. 21. Dorsal mandible ; natural size.
- Fig. 22. Dorsal aspect of the shell ; natural size.
- Fig. 23. Lateral aspect of the shell ; natural size.
- Fig. 24. Ventral aspect of the shell ; natural size.