

PLATE XXI.

Figs. 1-10. *SEPIA (METASEPIA) PFEFFERI*, n. sp., . . .

PAGE

145

- Fig. 1. Dorsal view of the specimen ; natural size.
- Fig. 2. Ventral view of the anterior portion ; natural size.
- Fig. 3. Circumoral region, showing the lips, buccal membrane, spermatic pad with some spermatophores upon it, and the bases of the arms ; magnified 2 diameters.
- Fig. 4. Dorsal aspect of the shell ; natural size.
- Fig. 5. Ventral aspect of the shell ; natural size.
- Fig. 6. Inner aspect of one of the arms ; natural size.
- Fig. 7. The left tentacular club ; magnified about 3 diameters.
- Fig. 8. The ventral mandible ; natural size.
- Fig. 9. The dorsal mandible ; natural size.
- Fig. 10. A sucker from one of the sessile arms ; magnified 15 diameters.

Figs. 11-19. *SEPIA ANDREANOIDES*, n. sp., . . .

139

- Fig. 11. Ventral view of a specimen, with the tentacles retracted ; natural size.
- Fig. 12. Dorsal view, with extended tentacles ; natural size.
- Fig. 13. The right tentacular club ; magnified about 3 diameters.
- Fig. 14. Dorsal aspect of the shell ; natural size.
- Fig. 15. Ventral aspect of the shell ; natural size.
- Fig. 16. Lateral aspect of the shell ; natural size.
- Fig. 17. A sucker from one of the sessile arms ; magnified 12 diameters.
- Fig. 18. Ventral mandible ; natural size.
- Fig. 19. Dorsal mandible ; natural size.