PLATE XXXIII.

		Figs. 1-11. Taonius hyperboreus, Steenstrup,	PAGE 191
Fig.	1.	Dorsal view of the type specimen; one-half natural size.	
Fig.	2.	Front view of an enlarged sucker, from one of the lateral arms; magnified 4 diameters.	
Fig.	3.	Side view of the same sucker; magnified 4 diameters.	
Fig.	4.	Front view of a similar sucker; less enlarged.	
Fig.	5.	Side view of the same sucker as fig. 4; less enlarged.	
Fig.	6.	The inner aspect of the tentacular club; natural size.	
Fig.	7.	A portion of the stem; enlarged to show the suckers and fixing- cushions.	
Fig.	8.	Three views of a large tentacular sucker; somewhat magnified.	
Fig.	9.	Front view of a similar sucker; magnified about 10 diameters.	
Fig. 1	0.	A sucker and fixing-cushion from the stem of the tentacle; magnified about 30 diameters.	
Fig. 1	1.	The gladius of a smaller specimen from the "Porcupine" collection; natural size.	
Fi Museur	_	es 1, 4, 5, 6, 7, 8, were drawn by Mr. Thornam from the type specimen in the Copenhagen	e
		Figs. 12-15. Calliteuthis reversa, Verrill,	183
Fig. 1	2.	The right tentacular club; magnified about 5 diameters.	
Fig. 1	3.	One of the larger tentacular suckers; magnified 40 diameters.	
Fig. 1	4.	A portion of the horny ring and papillary area; magnified 250 diameters.	14
Fig. 1	l 5.	Dorsal and lateral views of the gladius; natural size.	