

*Pilumnus granulatus*, Krauss. Natal. (The *Pilumnus xanthoides*, Krauss, from Natal, which is included by Dr. Kossmann in this genus, differs from the typical *Pilumni* in the form of the antero-lateral marginal teeth.)

*West Atlantic Species.*

*Pilumnus aculeatus* (Say). East coast of United States; West Indies (36 fathoms, A. Milne Edwards).

*Pilumnus gemmatus*, Stimpson. West Indies; Florida (to 37 fathoms). In some particulars this species approaches the genus *Heteractæa*.

*Pilumnus reticulatus*, Stimpson. St. Thomas.

*Pilumnus floridanus*, Stimpson. Tortugas; Brazil, off Barra Grande, 30 to 350 fathoms; Bahia (?).

*Pilumnus lacteus*, Stimpson. West Indies (to 37 fathoms, A. Milne Edwards).

*Pilumnus nudifrons*, Stimpson. West Indies; Sombrero Key (to 125 fathoms); West Barbados (to 133 fathoms, A. Milne Edwards).

*Pilumnus granulimanus*, Stimpson. Cuba.

*Pilumnus melanecanthus*, Kingsley. Florida.

*Pilumnus dasypodus*, Kingsley. Florida.

*Pilumnus caribæus*, Desbonne and Schramm. Guadeloupe.

*Pilumnus vinaceus*, A. Milne Edwards. Florida (to 37 fathoms, A. Milne Edwards). Martinique.

*Pilumnus gracilipes*, A. Milne Edwards. Barbados, 100 fathoms (A. Milne Edwards).

*Pilumnus quoyi*, Milne Edwards. Rio de Janeiro; Guiana.

*Pilumnus urinator*, A. Milne Edwards. Santa Cruz, 245 fathoms.

*Pilumnus miersii*, A. Milne Edwards. West Indies.

*Pilumnus tessellatus*, A. Milne Edwards. Brazil, Desterro.

*Pilumnus fragosus*, A. Milne Edwards. St. Thomas; Bahia, 7 to 20 fathoms.

*Pilumnus brasiliensis*, n. sp. Bahia, 7 to 20 fathoms.

*West American Species.*

*Pilumnus xantusii*, Stimpson. California, Cape St. Lucas.

*Pilumnus depressus*, Stimpson. California, Cape St. Lucas.

*Pilumnus stimpsonii* = *Pilumnus marginatus*, Stimpson.<sup>1</sup> California, Cape St. Lucas.

*Pilumnus spinohirsutus* (Lockington). California.

*Pilumnus limosus*, Smith. Peru; Panama.

<sup>1</sup> The name *marginatus* is preoccupied by Stimpson himself, for an Oriental species.