

The following species, besides those referred to below, are probably well established:—

Section A:—

*Matuta lunaris* (Herbst) = *Matuta planipes*, Fabricius (*vide* Hilgendorf), and *Matuta rubrolineata*, Miers, *Matuta leneifera*, Miers, var. Indian and Indo-Malaysian Seas; Pacific Ocean; Chefoo, Australia.

*Matuta circulifera*, Miers. Indo-Malaysian Seas. (Regarded by M. de Man as a variety of *Matuta lunaris*.)

Section B:—

*Matuta granulosa*, Miers. Oriental Seas; Amboina.

*Matuta maculata*, Miers. Chinese Seas; Panagatan Shoal.

*Matuta picta*, Hess (= *Matuta distinguenda*, Hoffman, and *Matuta obtusifrons*, Miers, var. ?). Indo-Pacific Seas.

*Matuta victrix*, Fabricius.

*Cancer victor*, Fabricius, Species Insect., ii. (Appendix) p. 502, 1781.

*Matuta victor*, Fabricius, Entom. Syst. Suppl., p. 369, 1798.

„ Milne Edwards, Hist. Nat. Crust., vol. ii. p. 115, pl. xx. figs. 3–6, 1837.

„ Hilgendorf, Monatsber. d. k. preuss. Akad. d. Wiss. Berlin, p. 810, 1878.

*victrix*, Miers, Trans. Linn. Soc. Lond. (Zool.), ser. 2, vol. i. pt. v. p. 243, pl. xxxix. figs. 1–3, 1877, et synonyma.

„ „ de Man, Notes Leyden Mus., iii. p. 110, 1881.

Philippine Islands, Samboangan, 10 to 20 fathoms. Two males, and an adult and a sterile female.

Adult ♂.	Lines.	Millims.
Length of carapace, . . . . .	19	40·5
Breadth of carapace, about . . . . .	20½	43

*Matuta victrix*, var. *crebrepunctata*, Miers.

*Matuta victrix*, var. *crebrepunctata*, Miers, *tom. cit.*, p. 244, pl. xxxix. fig. 4, 1877, not synonym.

Ternate, October 15, 1874 (an adult male).

Adult ♂.	Lines.	Millims.
Length and breadth of carapace, about . . . . .	15	31·5

*Matuta banksii*, Leach.

*Matuta banksii*, Leach, Zool. Miscell., vol. iii. p. 14, 1817 (?).

„ „ Miers, *tom. cit.*, p. 245, pl. xl. figs. 1, 2, 1877.

„ „ de Man, *tom. cit.*, p. 115, 1881.

Samboangan, Philippine Islands, 10 fathoms (two males and a female, with *Matuta victrix*).