

## PLATE VIII.

	Diam.	Page
Fig. 1. <i>Notolopas brasiliensis</i> , n. sp., male example, . . . . .	x 3	64
Fig. 1a. Post-abdomen of the male, . . . . .	x 8	
Fig. 1b. Inferior view of the cervical region of a variety with more strongly divergent rostral spines, showing also the form of the basal antennal joint and exterior maxillipedes in this species (from an adult male), . . . . .	x 5½	
Fig. 1c. <i>Notolopas lamellatus</i> , Stimpson (?), adult male; inferior view of the cervical region, showing the form of the basal antennal joint and exterior maxillipedes (figured for comparison from a specimen in the collection of the British Museum), . . . . .	x 5	
Fig. 2. <i>Micippa spinosa</i> , Stimpson, adult male, . . . . .	x 2	70
Fig. 2a. Rostrum of the same species; anterior view, . . . . .	x 2½	
Fig. 2b. Chelipede, . . . . .	x 3½	
Fig. 3. <i>Micippa spinosa</i> , var. <i>affinis</i> , Miers, adult male, . . . . .	x 2	70
Fig. 3a. Rostrum; anterior view, . . . . .	x 3	
Fig. 3b. Chelipede, . . . . .	x 5	
Fig. 3c. Post-abdomen of the male . . . . .	x 5	