

St. Paul's Rocks, 10 to 80 fathoms (and shore?).

<i>Stenorhynchus (?) spinifer</i> , n. sp.		<i>Grapsus maculatus</i> (Catesby).
--	--	-------------------------------------

Tristan da Cunha Group, Nightingale Island, 100 fathoms.

<i>Pilumnoplax heterochir</i> , Studer.

Cape of Good Hope, Simon's Bay, shallow water (5 to 18 fathoms).

<i>Stenorhynchus falcifer</i> , Stimpson.		<i>Pericera cornuta</i> , Milne Edwards.
<i>Achæopsis spinulosus</i> , Stimpson.		<i>Plagusia chabrus</i> (Linné).
<i>Dehaanius dentatus</i> (Milne Edwards).		<i>Hymenosoma orbiculare</i> (Desmarest).
		<i>Calappa flammea</i> (Herbst).
		<i>Mursia cristimana</i> (Desmarest).

South Africa, Wellington and Cape Town (rivers).

<i>Thelphusa (Potamonautes) perlata</i> , Milne Edwards.
--

Sea Point.

<i>Mursia cristimana</i> (Desmarest).

STATION 142.—Agulhas Bank, 150 fathoms; December 18, 1873; lat. $35^{\circ} 4' 0''$ S., long. $18^{\circ} 37' 0''$ E.; bottom, green sand; bottom temperature, $47^{\circ} 0$. Dredged.

<i>Lispognathus thomsoni</i> (Norman).		<i>Litocheira kingsleyi</i> , n. sp.
<i>Pilumnoplax heterochir</i> , Studer.		<i>Mursia cristimana</i> (Desmarest).
<i>Ebalia tuberculosa</i> (A. Milne Edwards) (yg. ?).		

INDO-PACIFIC OR ORIENTAL REGION.

STATION 161.—Port Philip, 33 fathoms; April 1, 1874; lat. $38^{\circ} 22' 30''$ S., long. $144^{\circ} 36' 30''$ E.; bottom, sand. Trawled.

<i>Achæus tenuicollis</i> , n. sp.		<i>Portunus corrugatus</i> (Pennant).
<i>Pilumnus tomentosus</i> , Latreille.		<i>Halicarcinus ovatus</i> , Stimpson.
<i>Ebalia (Phlyxia) intermedia</i> , n. sp.		