

STATION 233B.—Japan, 15 fathoms; May 26, 1875; lat. $34^{\circ} 18' 0''$ N., long. $133^{\circ} 35' 0''$ E.; bottom, blue mud. Trawled.

<i>Neptunus (Amphitrite) hastatoides</i> , (Fabricius.)	<i>Pilumnoplax vestita</i> (de Haan), var. <i>sexdentata</i> , Haswell.
<i>Arcania septemspinosa</i> (Fabricius).	

Japan, near Lake Biva.

Thelphusa (Geothelphusa) dehaanii, White.

Japan, Hakouni (2500 feet).

Thelphusa (Geothelphusa) dehaanii, White.

STATION 237.—Japan, near Yokohama, 1875 fathoms; June 17, 1875; lat. $34^{\circ} 37' 0''$ N., long. $140^{\circ} 32' 0''$ E.; bottom, blue mud; bottom temperature, $35^{\circ}\cdot 3$. Trawled.

Ethusa (Ethusina) challengerii, n. sp.

Sandwich Islands, Hawaii.

Neptunus (Neptunus) sanguinolentus (Herbst).

Sandwich Islands, Hilo (beach).

Metopograpsus messor (Forskål).

Sandwich Islands, Honolulu (reefs), some 16, 18 to 20 fathoms.

<i>Carpilius maculatus</i> (Linné).	<i>Neptunus (Neptunus) sanguinolentus</i> (Herbst).
<i>Lophactæa granulosa</i> (Rüppell).	
<i>Actæa nodulosa</i> , White.	<i>Thalamita integra</i> , Dana.
<i>Etisus lærivimanus</i> , Randall.	<i>Podophthalmus vigil</i> (Fabricius).
<i>Trapezia rufopunctata</i> , var. <i>guttata</i> , Rüppell.	<i>Calappa hepatica</i> (Linné).

Society Islands, Tahiti, Papiete.

<i>Scylla serrata</i> (Forskål).	<i>Gelasimus tetragonon</i> (Herbst).
<i>Cardiosoma carnifex</i> (Herbst).	<i>Metopograpsus messor</i> (Forskål).