

The Family here contains :—

1. *Crisia.*

- (1) *Crisia biciliata*, Macgilliv. (Pl. I. figs. 1, 2).
- (2) *Crisia eburnea*, var. *laxa* (Pl. II. fig. 1).
- (3) *Crisia denticulata*, Lamk. (Pl. II. fig. 3).
Crisia denticulata, var. α , *gracilis* (Pl. I. fig. 4).
Crisia denticulata, var. β , *patagonica*.
- (4) *Crisia elongata*, M.-Edw. (Pl. I. fig. 3).
- (5) *Crisia acuminata*, n. sp. (Pl. III. fig. 1).
- (6) *Crisia acropora*, Bk.
- (7) *Crisia holdsworthii*, Bk. (Pl. III. fig. 2).
- (8) *Crisia conferta*, Bk. (Pl. II. fig. 5).
- (9) *Crisia cylindrica*, n. sp. (Pl. II. figs. 2, 4).

1. *Crisia.*

- Sertularia* (pars), Linné.
- Cellularia* (pars), Pallas.
- Cellaria* (pars), Solander, Lamk.
- Tibiana* (pars), Lamk., Schweigger, Blainville, Lister.
- Falcaria* (pars), Oken, Gray.
- Eucratea* (pars), Esper, Lamx., Fleming, &c.
- Crisia* (pars), Lamx., Johnst., Gray, Auctt.
- Crisidia* (pars), M.-Edwards, Busk, d'Orbigny (1852).
- Filicrisia*, d'Orbigny (1851).

Character.—Zoœcia disposed in a single or double longitudinal series. Oœcia modified zoœcia, with a tubular aperture, walls punctate.

§ a. Uniserial.

- Crisidia*, M.-Edwards, Busk, Auctt.
- Unicellaria* (pars), Blainville.
- Falcaria*, Oken, Gray.
- Tibiana*, Lister (*nec* Blainville, Lamx.).
- Unicrisia*, d'Orb., Vine.

No species belonging to this section occurs in the Challenger collection.

§ β. Zoœcia disposed in a double series, opposite or alternate.

§§ a. Zoœcia opposite in pairs, from two to four (rarely six) in an internode.

- Bicrisia*, d'Orbigny.
- Crisia* (pars), Auctt.