Genus Marginaster, Perrier.

Marginaster, Perrier, Bull. Mus. Comp. Zoöl. Harvard, 1881, vol. ix. No. 1, p. 16; Nouv. Archives Mus. Hist. Nat., 2e Série, 1884, t. vi. p. 229.

Two species bearing this generic name were described by Perrier in 1881, and a third species was added by the same author in the following year. A short diagnosis of the genus was first given in 1884. The descriptions of these interesting forms are unfortunately so brief and general that although two of the species are figured, I feel some doubt as to whether all the three belong to one genus, and I have been unable to form any clear view as to the scope of the genus. It is, therefore, not without some hesitation that I now venture to describe a small starfish dredged during the "Porcupine" Expedition as a new species of Marginaster, but its characters appear to me to accord so closely with those mentioned as presented by Marginaster pectinatus, Perrier (which I presume may be considered as the type of the genus), that I prefer to take this course rather than burden nomenclature with what might ultimately prove to be a superfluous generic name. Under these circumstances it would be premature and injudicious in me to endeavour to formulate a more complete diagnosis of the genus, on the basis of the form which I have studied, until fuller information on the previously described species is available.

Chorology of the Genus Marginaster.

a. Geographical distribution:—

ATLANTIC: Four species between the parallels of 10° and 60° N.

Marginaster pectinatus, in the Gulf of Mexico, off the coast of Yucatan. Marginaster echinulatus, off Barbados. Marginaster pentagonus, "Talisman" dredging No. 37. Marginaster fimbriatus, between the north of Ireland and Rockall.

- β. Bathymetrical range: 69 to 1360 fathoms.
- γ. Nature of the Sea-bottom: Not recorded.

Chorological Synopsis of the Species.

| Marginaster echinulatus | Ocean. Atlantic. | Range in Fathoms. | Nature of the Sca-bottom. | |
|--------------------------|------------------|-------------------|---------------------------|------|
| | | | | 1111 |
| Marginaster fimbriatus . | Atlantic. | 1360 | ••• | |
| Marginaster pentagonus . | Atlantic. | 218 | ••• | ••• |
| Marignaster pectinatus . | Atlantic. | 95 | | ••• |

¹ Rapport sur les Travaux de la Commission chargée par M. le Ministre de l'Instruction publique d'étudier la Faune sous-marine dans les grandes profondeurs de la Méditerranée et de l'Ocean Atlantique, par M. Alphonse Milne-Edwards, Membre de l'Institut. (Extrait des Archives des Missions scientifiques et littéraires. Troisième série.—Tome neuvième, Paris, 1882, p. 51.)

² Nouv. Archives Mus. Hist. Nat., 2e Série, 1884, t. vi. pl. i.