Patiria bellula. Shallow water.

Asterina exigua. Shallow water. Also off Samboangan, Philippine Islands; Sea Point, near Cape Town; and Port Jackson, Australia.

Stichaster felipes. Also at Station 142.

Cribrella ornata. Shallow water to 20 fathoms.

Asterias (Stolasterias) africana. Shallow water to 20 fathoms.

Sea Point, near Cape Town.

Asterina exigua. Also off Samboangan, Philippine Islands; Simon's Bay, Cape of Good Hope; and Port Jackson, Australia.

STATION 142. South of the Cape of Good Hope. December 18, 1873. Lat. 35° 4′ 0″ S., long. 18° 37′ 0″ E. Depth 150 fathoms. Green sand. Bottom temperature 47° 0 Fahr.; surface temperature 65° 5 Fahr.

Stichaster felipes. Also at Simon's Bay, Cape of Good Hope.

STATION 143. Off the Agulhas Bank, south of the Cape of Good Hope. December 19, 1873. Lat. 36° 48′ 0″ S., long. 19° 24′ 0″ E. Depth 1900 fathoms. Globigerina ooze. Bottom temperature 35° 6 Fahr.; surface temperature 73° 0 Fahr.

Pararchaster pedicifer. Also at Station 147.

STATION 145. Off Marion Island. December 27, 1873. Lat. 46° 43′ 0″ S., long. 38° 4′ 30″ E. Depth 140 fathoms. Volcanic sand. Surface temperature 41° 0 Fahr.

? Crossaster penicillatus. Also at Station 135c.

Pedicellaster hypernotius.

Off Marion Island. Depth 50 fathoms.

Leptoptychaster kerguelenensis. Also at Stations 149A, 149D, 149E. Gnathaster meridionalis. Also at Stations 149D, 149H, 150, and 151.

Gnathaster elongatus. Also at Stations 149H, 150, and 151.

Porania antarctica. Also off Prince Edward Island, and at Station 147.

Pteraster semireticulatus.

Cribrella simplex. Also off Inaccessible Island, and at Stations 1350, 145A, and 148.