Porania glaber. Also at Station 149E.
Retaster peregrinator.
Perknaster densus.
Echinaster spinulifer. Also at Station 149D.
Asterias meridionalis. Also at Stations 149D, 149E, and off Marion Island.
Labidiaster annulatus. Also at Stations 150, 151, and 191.

Off Kerguelen Island. Position not recorded.

Leptoptychaster kerguelenensis. Depth 10 to 50 and 100 fathoms. Also at Stations 149A, 149D, 149E, and off Marion Island.

Cribrella simplex, var. granulosa. Depth 10 to 50 fathoms. Also at Station 149D.

Asterias meridionalis. Depth 10 to 50 fathoms. Also at Stations 149D, 149E, 149H, and off Marion Island.

Asterias perrieri. Depth 110 fathoms. Also at Station 149D.

STATION 150. Between Kerguelen Island and Heard Island. February 2, 1874. Lat. 52° 4′ 0″ S., long. 71° 22′ 0″ E. Depth 150 fathoms. Coarse gravel. Bottom temperature 35° 2 Fahr.; surface temperature 37° 5 Fahr.

> Gnathaster meridionalis. Also at Stations 149D, 149E, 151, and off Marion Island.

> Gnathaster elongatus. Also at Stations 1495, 151, and off Marion Island.

Porania spiculata. Also at Stations 151 and 191.

Solaster subarcuatus.

Pteraster rugatus.

Asterias (Smilasterias) triremis.

Labidiaster annulatus. Also at Stations 149H, 151 and 191.

STATION 151. Off Heard Island. February 7, 1874. Lat. 52° 59' 30" S., long. 73° 33' 30" E. Depth 75 fathoms. Volcanic mud. Surface temperature 36° 2 Fahr. Bathybiaster loripes, var. obesa. Also at Station 149H. Gnathaster meridionalis. Also at Stations 149D, 149H, 150, and off Marion Island. Gnathaster elongatus. Also at Stations 149H, 150, and off Marion Island.