- STATION 55 (1869). In the Faeröe Channel. Lat. 60° 4' N., long. 6° 19' W. Depth 605 fathoms. Bottom temperature - 1°.2 C.; surface temperature 11°.4 C. Lophaster furcifer. Also at Station 52.
- STATION 57 (1869). In the Faeroe Channel. Lat. 60° 14' N., long. 6° 17' W. Depth 632 fathoms. Bottom temperature -0°.8 C.; surface temperature 11°.1 C. Plutonaster bifrons. Also at Stations 19 and 58. Ctenodiscus corniculatus. Also at Stations 58 and 82.

Crossaster papposus. Also at Stations 52 and 64.

Korethraster hispidus.

STATION 58 (1869). In the Faeröe Channel. Lat. 60° 21' N., long. 6° 51' W. Depth 540 fathoms. Bottom temperature -0°.6 C.; surface temperature 10°.6 C. Plutonaster bifrons. Also at Stations 19 and 57. Ctenodiscus corniculatus. Also at Stations 57 and 82.

In the Faeröe Channel. Lat. 60° 21' N., long. 5° 41' W. STATION 59 (1869). Depth 580 fathoms. Bottom temperature - 1°-3 C.; surface temperature 11°.5 C.

Hymenaster pellucidus.

- STATION 62 (1869). In the Faeroe Channel. Lat. 61° 59' N., long. 4° 38' W. Depth 125 fathoms. Bottom temperature 7°0 C.; surface temperature 9°8 C. Cribrella oculata. Also at Stations 51, 52, 54, 65, 74, and 90.
- STATION 64 (1869). Between the Faeröe and Shetland Islands. Lat. 61° 21' N., long. 3° 44' W. Depth 640 fathoms. Bottom temperature -1°1 C.; surface temperature 9°.3 C.

Crossaster papposus. Also at Stations 52 and 57.

STATION 65 (1869). In the Faeröe Channel. Lat. 61° 10' N., long. 2° 21' W. Depth 345 fathoms. Bottom temperature - 1°·1 C.; surface temperature 11°·1 C.

> Pontaster tenuispinus. Also at Stations 76 and 90. Leptoptychaster arcticus. Also at Station 82; and in 1870 at Station 3. Cribrella oculata. Also at Stations 51, 52, 54, 62, 74, and 90.