

- STATION 67 (1869). East of the Shetland Islands. Lat. $60^{\circ} 32' N.$, long. $0^{\circ} 29' W.$ Depth 64 fathoms. Bottom temperature $9^{\circ} 5' C.$; surface temperature $11^{\circ} 0' C.$
Astropecten irregularis. Also at Stations 18, 46, and 68.
Luidia sarsii. Also at Stations 46 and 68.
Asterias (Leptasterias) mulleri. Also at Stations 68 and 82.
- STATION 68 (1869). East of the Shetland Islands. Lat. $60^{\circ} 23' N.$, long. $0^{\circ} 33' E.$ Depth 75 fathoms. Bottom temperature $6^{\circ} 7' C.$; surface temperature $11^{\circ} 4' C.$
Astropecten irregularis. Also at Stations 18, 46, and 67.
Luidia sarsii. Also at Stations 46 and 67.
Asterias (Leptasterias) mulleri. Also at Stations 67 and 82.
- STATION 74 (1869). East of the Shetland Islands. Lat. $60^{\circ} 39' N.$, long. $3^{\circ} 9' W.$ Depth 203 fathoms. Bottom temperature $8^{\circ} 7' C.$; surface temperature $11^{\circ} 4' C.$
Cribrella oculata. Also at Stations 51, 52, 54, 62, 65, and 90.
Brisinga endecacnemos. Also at Station 45, and off Valentia; and in 1870 at Stations 4, 6, 14, and 17.
- STATION 76 (1869). In the Faerøe Channel. Lat. $60^{\circ} 36' N.$, long. $3^{\circ} 58' W.$ Depth 344 fathoms. Bottom temperature $-1^{\circ} 1' C.$; surface temperature $10^{\circ} 1' C.$
Pontaster tenuispinus. Also at Stations 65 and 90.
Psilaster andromeda. Also in 1870 at Station 3.
Bathybiaster vexillifer.
- STATION 82 (1869). In the Faerøe Channel. Lat. $60^{\circ} 0' N.$, long. $5^{\circ} 13' W.$ Depth 312 fathoms. Bottom temperature $5^{\circ} 2' C.$; surface temperature $11^{\circ} 2' C.$
Ctenodiscus corniculatus. Also at Stations 57 and 58.
Leptoptychaster arcticus. Also at Station 65; and in 1870 at Station 3.
Asterias (Leptasterias) mulleri. Also at Stations 67 and 68.
- STATION 87 (1869). North-west of the Hebrides. Lat. $59^{\circ} 35' N.$, long. $9^{\circ} 11' W.$ Depth 767 fathoms. Bottom temperature $5^{\circ} 2' C.$; surface temperature $11^{\circ} 4' C.$
Zoroaster fulgens.
- STATION 90 (1869). In the Faerøe Channel. Lat. $59^{\circ} 41' N.$, long. $7^{\circ} 34' W.$ Depth 458 fathoms. Bottom temperature $7^{\circ} 3' C.$; surface temperature $11^{\circ} 7' C.$
Pontaster tenuispinus. Also at Stations 65 and 76.
Cribrella oculata. Also at Stations 51, 52, 54, 62, 65, and 74.