

STATION 10. In the Faerøe Channel. August 24, 1882. Lat. $59^{\circ} 40' 0''$ N., long. $7^{\circ} 21' 0''$ W. Depth 516 fathoms. Bottom temperature $46^{\circ} 0$ Fahr.

Plutonaster bifrons. Also at Station 11.

Psilaster andromeda. Also at Station 11.

Mimaster tizardi. Also at Station 11.

Cribrella oculata, var. (see p. 543). Also at Stations 1 and 11.

STATION 11. In the Faerøe Channel. August 28, 1882. Lat. $59^{\circ} 29' 0''$ N., long. $7^{\circ} 13' 0''$ W. Depth 555 fathoms. Bottom temperature $45^{\circ} 5$ Fabr.

Plutonaster bifrons. Also at Station 10.

Psilaster andromeda. Also at Station 10.

Mimaster tizardi. Also at Station 10.

Zoroaster fulgens. Also at Station 13.

Cribrella oculata, var. (see p. 543). Also at Stations 1 and 10.

STATION 13. In the Faerøe Channel. August 31, 1882. Lat. $59^{\circ} 51' 2''$ N., long. $8^{\circ} 18' 0''$ W. Depth 570 fathoms. Bottom temperature $45^{\circ} 7$ Fabr.

Zoroaster fulgens. Also at Station 11.

V. "LIGHTNING" EXPEDITION.

STATION 6. In the Faerøe Channel. Lat. $60^{\circ} 45' 0''$ N., long. $4^{\circ} 49' 0''$ W. Depth 510 fathoms. Bottom temperature $0^{\circ} 5$ C.

Pontaster tenuispinus.

STATION 7. In the Faerøe Channel. September 3, 1868. Lat. $60^{\circ} 7' 0''$ N., long. $5^{\circ} 21' 0''$ W. Depth 500 fathoms. Bottom temperature $1^{\circ} 1$ C.; surface temperature $10^{\circ} 5$ C.

Odinia pandina.

Off Valentia. Depth and conditions not stated.

Porania pulvillus.