

III. "KNIGHT ERRANT" EXPEDITION.

(1.) LITTORAL ZONE.

Species found in depths which extend from the Shore to 150 fathoms.

	Depth in Fathoms.	Occurrence in other Zones.
<i>ASTROPECTINIDÆ.</i>		
<i>LUIDIINÆ.</i>		
<i>Luidia sarsii</i>	53
<i>GYMNASTERIIDÆ.</i>		
<i>Porania pulvillus</i>	53
<i>SOLASTERIDÆ.</i>		
<i>SOLASTERINÆ.</i>		
<i>Crossaster papposus</i>	53
<i>ECHINASTERIDÆ.</i>		
<i>ECHINASTERINÆ.</i>		
<i>Gribrella oculata</i>	53
<i>ASTERIIDÆ.</i>		
<i>Asterias rubens</i>	53
<i>Asterias (Leptasterias) mülleri</i>	53

(2.) CONTINENTAL ZONE.

Crossaster papposus, var. septentrionalis 375 fathoms.

(3.) ABYSSAL ZONE.

Species found in depths greater than 500 fathoms.

	At or greater than						Greatest Depth recorded.	Occurrence in other Zones.
	500	750	1000	1500	2000	2500		
<i>ARCHASTERIDÆ.</i>								
<i>PABAROHASTERINÆ.</i>								
<i>Pontaster tenuispinus, var. platynota</i>	+	540
<i>PLUTONASTERINÆ.</i>								
<i>Plutonaster bifrons</i>	+	540
<i>ASTROPECTINIDÆ.</i>								
<i>ASTROPECTININÆ.</i>								
<i>Psilaster andromeda</i>	+	530
<i>PENTAGONASTERIDÆ.</i>								
<i>MIMASTERINÆ.</i>								
<i>Minaster tizardi</i>	+	555