

IV. "TRITON" EXPEDITION.

(1.) LITTORAL ZONE.

Species found in depths which extend from the Shore to 150 fathoms.

	Depth in Fathoms.	Occurrence in other Zones.
<i>ASTROPECTINIDÆ.</i>		
<i>LUIDIINÆ.</i>		
<i>Luidia ciliaris</i>	87
<i>ANTHENEIDÆ.</i>		
<i>Hippasteria plana</i>	87

(2.) CONTINENTAL ZONE.

Species found in depths between 150 fathoms and 500 fathoms.

	Depth in Fathoms.	Occurrence in other Zones.
<i>GYMNASTERIIDÆ.</i>		
<i>Rhögaster murrayi</i>	433
<i>ECHINASTERIDÆ.</i>		
<i>ECHINASTERINÆ.</i>		
<i>Cribrella oculata</i>	240	Also Abyssal.
<i>ASTERIIDÆ.</i>		
<i>Asterias (Leptasterias) mülleri</i>	433

(3.) ABYSSAL ZONE.

Species found in depths greater than 500 fathoms.

	At or greater than						Greatest Depth recorded.	Occurrence in other Zones.
	500	750	1000	1500	2000	2500		
<i>ARCHASTERIDÆ.</i>								
<i>PARARHASTERINÆ.</i>								
<i>Pontaster tenuispinus</i> , var. <i>platynota</i>	+	608
<i>PLUTONASTERINÆ.</i>								
<i>Plutonaster bifrons</i>	+	555
<i>ASTROPECTINIDÆ.</i>								
<i>ASTROPECTININÆ.</i>								
<i>Psilaster andromeda</i>	+	555
<i>PENTAGONASTERIDÆ.</i>								
<i>MIMASTERINÆ.</i>								
<i>Mimaster tizardi</i>	+	555
<i>ZOROASTERIDÆ.</i>								
<i>Zoroaster fulgens</i>	+	570
<i>PTERASTERIDÆ.</i>								
<i>PTERASTERINÆ.</i>								
<i>Pteraster militaris</i>	+	530
<i>Pteraster militaris</i> , var. <i>prolata</i>	+	608
<i>ECHINASTERIDÆ.</i>								
<i>ECHINASTERINÆ.</i>								
<i>Cribrella oculata</i> , var. (see p. 543)	+	555	Also Continental.