

EUASTEROIDEA), WITH THE PRINCIPAL LOCALITIES, THE GENERAL DISTRIBUTION, AND THE SYNONYMS.

Previously known species collected by the Challenger are marked \*.

L	GEOGRAPHICAL AREAS.							BATHYMETRICAL ZONES.			BOTTOM TEMPERATURE: FAHRENHEIT.	SYNONYMA AND REMARKS.	
	I. North Atlantic.	II. South Atlantic.	III. Indian Ocean.	IV. Southern Ocean.	V. Eastern Archipago.	VI. North Pacific.	VII. South Pacific.	L	Littoral, 0-150 Fathoms.	Continental, 150-500 Fathoms.	Absyal, greater than 500 Fathoms.		
								C.	C.	A.	A.		
(1)	+	...	...	...	...	...	...	...	A	37°·2-38°·0	.....		
(2)	...	...	...	+	...	...	...	...	A	...	...	.....	
(3)	...	...	...	+	...	...	...	...	A	34°·2-35°·6	.....		
(4)	...	...	...	...	...	+	...	...	A	35°·3-38°·1	.....		
(5)	+	...	...	...	...	...	...	...	A	36°·2-37°·2	.....		
(6)	+	...	...	...	...	...	...	...	A	40°·0	= <i>Archaster simplex</i> , Perrier.		
(7)	...	+	...	...	...	...	...	...	C	40°·3	.....		
(8)	+	...	...	...	...	...	...	...	A	36°·2-38°·0	.....		
(9)	...	...	...	+	...	...	...	...	A	35°·6	.....		