

		LOCALITIES.	RANGE IN DEPTH: FATHOMS.	NATURE OF THE SEA- BOTTOM.
PHANEROZONIA.				
ASTROPECTINIDÆ.				
ASTROPECTININA.				
<i>Leptoptychaster</i> , Smith.				
* <i>L. antarcticus</i> , Sladen . . {	Between Marion and Kerguelen Islands.	210	{ Hard ground (Gravel, shells).	(1)
<i>L. arcticus</i> , Sars, sp. . . {	Off North Cape; coast of Norway; E. of N. America.	20-690	Clay, sometimes sandy.	(2)
* <i>L. arcticus</i> , var. <i>elongata</i> , Sladen . .	E. of N. America.	85-1350	Gravel, stones; Bluemud.	(3)
* <i>L. kerguelensis</i> , Smith . .	Kerguelen Isld.; Marion Isld.	10-100	Volcanic mud.	(4)
<i>Moiraster</i> , Sladen.				
<i>M. magnificus</i> , Bell, sp. . . .	St. Helena	?	?	(5)
<i>Astropecten</i> , Linck.				
* <i>A. acanthifera</i> , Sladen . . .	Banda Sea.	140	Blue mud.	(6)
<i>A. alatus</i> , Perrier . . .	?	?	?	(7)
<i>A. alligator</i> , Perrier . . .	Off Florida.	24 (?)	(8)
<i>A. andersoni</i> , Sladen . . .	Mergui Archipelago.	10	(9)
<i>A. antillensis</i> , Lütken . . .	St. Thomas; Guadeloupe.	Shallow.	(10)
<i>A. arenarius</i> , Perrier . . .	?	?	?	(11)
<i>A. articulatus</i> , Say . . .	From New Jersey to W. Indies.	Shallow.	(12)
<i>A. aurantiacus</i> (Linné), Gray {	Mediterranean; Adriatic; Madeira; Canary Islands.	10-100	(13)
<i>A. bipinnosus</i> (Otto), Müller and Troschel	Mediterranean; Adriatic.	2-35	(14)
* <i>A. brasiliensis</i> , Müller and Troschel	St. Vincent; Fernando Noronha; Brazil.	7-20	(15)