

	LOCALITIES.	RANGE IN DEPTH: FATHOMS.	NATURE OF THE SEA-BOTTOM.	
PHANEROZONIA.				
<i>PENTAGONASTERIDÆ.</i>				
<i>PENTAGONASTERINÆ.</i>				
<i>Ohitonaster</i> , Sladen.				
* <i>C. cataphractus</i> , Sladen	62° 26' 0" S., 95° 44' 0" E.	1976	Diatom ooze.	(1)
<i>Calliderma</i> , Gray.				
<i>C. emma</i> , ¹ Gray	Japan (?).	†	†	(2)
<i>Iconaster</i> , Sladen.				
<i>I. longimanus</i> , Mobius, sp.	{ Malacca (?); Percy Island (Queensland); Prince of Wales Channel (<i>vide</i> Bell). }	7	†	(3)
<i>Gnathaster</i> , Sladen.				
<i>G. dilatatus</i> , Perrier, sp.	New Zealand.	(4)
* <i>G. elongatus</i> , Sladen	{ Off Marion Island; Kerguelen Island; Heard Island. }	50-150	{ Volcanic mud; coarse gravel. }	(5)
<i>G. grayi</i> , Bell, sp.	{ Sandy Point (Strait of Magellan). }	9-10	Sand.	(6)
* <i>G. meridionalis</i> , Smith, sp.	{ Off Marion Island; Kerguelen Island; Heard Island. }	5-150	{ Volcanic mud; coarse gravel. }	(7)
<i>G. miliaris</i> , Gray, sp.	New Zealand.	(8)
<i>G. parillosus</i> , Gray, sp.	{ N. Australia; Sandy Point (Strait of Magellan). }	9-10	Sand.	(9)
* <i>G. pilulatus</i> , Sladen	{ Cape Virgins, off Strait of Magellan. }	55	Sand.	(10)
<i>G. singularis</i> , Muller and Troschel, sp.	{ W. of South America; Tom Bay (<i>vide</i> Bell). }	{ Shallow to 30 }	Rock; kelp; mud	(11)
<i>G. (?) verrucosus</i> , Philippi, sp.	Near Valparaiso.	(12)
<i>Nymphaster</i> , Sladen.				
* <i>N. albidus</i> , Sladen	Off Cape Verde Islands.	(13)