

GEOGRAPHICAL AREAS.							BATYMETRICAL ZONES.			BOTTOM TEMPERATURE: FAHRENHEIT.			SYNONYMA AND REMARKS.
I.	II.	III.	IV.	V.	VI.	VII.	Littoral, 0-150 Fathoms.	Continental, 150-500 Fathoms.	Absyal, greater than 500 Fathoms.	L	O	A	
(1)	+	L
(2)	L
(3)	+	...	L
(4)	?	L	= <i>Oreaster carinatus</i> , Müller and Troschel.
(5)	L	= <i>Oreaster chinensis</i> , Müller and Troschel.
(6)	L	Probably includes <i>Oreaster occidentalis</i> , Verrill.
(7)	+	...	L	= <i>Oreaster decipiens</i> , Bell.
(8)	+	L	{ = <i>Asterias dorsatus</i> , Linnaé; <i>Oreaster dorsatus</i> , Lütken. Incl.: <i>Asterias nodosa</i> (pars), Linnaé; <i>Oreaster olivatus</i> , Müller and Troschel.
(9)	+	L
(10)	L	= <i>Oreaster gracilis</i> , Lütken.
(11)	+	L	{ = <i>Oreaster granulosus</i> , Bell. (A doubtful member of the genus.)
(12)	+	...	+	...	L	{ = <i>Pentaceros nodosa</i> , Gray (non Linnaé); <i>Oreaster nodosus</i> , Müller and Troschel; <i>Oreaster grayi</i> , Bell. Incl.: <i>Oreaster Clouei</i> , Perrier. (Probably the same species as <i>Pentaceros verrucosus</i> , Müller and Troschel, sp., which has priority.)
(13)	+	...	L	= <i>Oreaster Hedemanni</i> , Lütken.
(14)	+	L	{ = <i>Pentaceros gibbus</i> , Linck; <i>Oreaster hilucus</i> , Müller and Troschel.
(15)	+	...	L	= <i>Oreaster luetkeni</i> , Bell.
(16)	+	L	{ = <i>Asterias mammillata</i> , Audouin; <i>Oreaster mammillatus</i> , Müller and Troschel.
(17)	+	...	+	...	L	{ = <i>Pentaceros gibbus</i> , muricatus, Linck; <i>Pentaceros muricatus</i> , Gray. Incl.: <i>Asterias Linkii</i> , de Blainville; <i>Oreaster muricatus</i> , Dujardin and Hupé; <i>Oreaster Linckii</i> , Lütken; <i>Oreaster castellum</i> , Grube; <i>Oreaster nodosus</i> , var. <i>muri-</i> catus, von Martens.