

	LOCALITIES.	RANGE IN DEPTH: FATHOMS.	NATURE OF THE SEA-BOTTOM.	
PHANEROZONIA.				
<i>PENTACEROTIDÆ.</i>				
<i>Pentaceros</i> , Linck.				
<i>P. muelleri</i> , Bell, sp.	Billiton.	(1)
<i>P. nodulosus</i> , Perrier.	W. and N.W. Australia.	(2)
<i>P. occidentalis</i> , Verrill, sp. {	Margarita Bay; Cape San Lucas; Panama; Guayaquil. }	12-18	Rocky ground.	(3)
<i>P. orientalis</i> , Müller and Troschel, sp. {	China; New Britain (<i>vide</i> Studer). }	(4)
<i>P. productus</i> , Bell, sp.	Billiton.	(5)
* <i>P. productus</i> , var. <i>tubercata</i> , Sladen	Off Philippine Islands.	10	Sand.	(6)
<i>P. regulus</i> (Val.), Müller and Troschel, sp. }	Pondicherry (Bay of Bengal).	(7)
<i>P. reinhardti</i> , Lütken, sp.	Nicobar Islands	(8)
<i>P. reticulatus</i> , Linck }	Florida; Bahamas; Barbados; Antilles; Brazil; Abrolhos reefs; Freemantle (West Australia). }	(9)
<i>P. sladeni</i> , de Loriol	Mauritius.	(10)
<i>P. superbus</i> , Möbius, sp.	Sumatra; Madras (<i>vide</i> Bell).	(11)
<i>P. thurstoni</i> , Bell, sp.	Tuticorin (Madras).	(12)
<i>P. troscheli</i> , Bell, sp.	Billiton.	(13)
<i>P. tuberculatus</i> , Müller and Troschel }	Red Sea; E. coast of Africa.	(14)
* <i>P. turritus</i> , Linck }	Indian Ocean generally; Eastern Archipelago; New Guinea; Port Essington; Cape Grenville; Port Denison; New Caledonia; New Britain. }	4-28	(15)