

	GEOGRAPHICAL AREAS.							BATHYMETRICAL ZONES.			BOTTOM TEMPERATURE: FAHRENHEIT.	SYNONYMA AND REMARKS.
	I. North Atlantic.	II. South Atlantic.	III. Indian Ocean.	IV. Southern Ocean.	V. Eastern Archipelago.	VI. North Pacific.	VII. South Pacific.	L. Littoral, 0-100 Fathoms.	O. Continental, 150-500 Fathoms.	A. Abyssal, greater than 500 Fathoms.		
(1)	+	L	= <i>Oreaster muelleri</i> , Bell.
(2)	+	L	= <i>Oreaster nodulosus</i> , Bell.
(3)	+	+	L	= <i>Oreaster occidentalis</i> , Verrill. (Probably the same species as <i>Pentaceros Cumingii</i> , Gray.)
(4)	+	+	L	= <i>Oreaster orientalis</i> , Müller and Troschel.
(5)	+	L	= <i>Oreaster productus</i> , Bell.
(6)	+	L
(7)	+	L	= <i>Oreaster regulus</i> , Müller and Troschel.
(8)	+	L	= <i>Oreaster Reinhardti</i> , Lutken.
(9)	+	+	+	L	= <i>Pentaceros gibbus</i> , <i>reticulatus</i> , Linck; <i>Asterias reticulata</i> , Linné; <i>Oreaster reticulatus</i> , Müller and Troschel. Incl: <i>Pentaceros lentiginosus</i> , Linck; <i>Asterias gigas</i> , Linné; <i>Asterias pentacyphus</i> , Retzius; <i>Asterias Schu.</i> , de Blainville; <i>Pentaceros grandis</i> , Gray; <i>Pentaceros gibbus</i> , Gray; <i>Pentaceros aculeatus</i> , Gray; <i>Oreaster aculeatus</i> , Müller and Troschel; <i>Oreaster lapidarius</i> , Grube; <i>Oreaster gigas</i> , Lütken; <i>Pentaceros tuberosus</i> (Behn, M.S.), Möbius.
(10)	+	L
(11)	+	...	+	L	= <i>Oreaster superbus</i> , Möbius.
(12)	+	L	= <i>Oreaster thurstoni</i> , Bell.
(13)	+	L	= <i>Oreaster troscheli</i> , Bell.
(14)	+	L	= <i>Oreaster tuberculatus</i> , Müller and Troschel. (Probably the same as <i>Pentaceros mammillatus</i> (Audouin), Parrier.)
(15)	+	...	+	...	+	L	= <i>Pentaceros gibbus</i> , <i>turritus</i> , Linck; <i>Oreaster turritus</i> , Müller and Troschel. Incl: <i>Asterias nodosa</i> (pars), Linné; <i>Pentaceros Franklinii</i> , Gray; <i>Pentaceros modestus</i> , Gray; <i>Oreaster mammosus</i> , Parrier; <i>Oreaster nodosus</i> , Bell.