

	GEOGRAPHICAL AREAS.							BATHYMETRICAL ZONES.				BOTTOM TEMPERATURE: FAHRENHEIT.	SYNONYMA AND REMARKS.
	I. North Atlantic.	II. South Atlantic.	III. Indian Ocean.	IV. Southern Ocean.	V. Eastern Archipelago.	VI. North Pacific.	VII. South Pacific.	Littoral, 0-150 Fathoms.	Continental, 150-600 Fathoms.	Abyssal, greater than 600 Fathoms.			
	I.	II.	III.	IV.	V.	VI.	VII.	L	O	A			
(1)	+	L		= <i>Oreaster valvatus</i> , Müller and Troschel.
(2)	?	L		{ <i>Oreaster verrucosus</i> , Müller and Troschel. Probably includes <i>Pentaceros nodosus</i> , Gray (non Linné), <i>Oreaster nodosus</i> , Müller and Troschel, <i>Oreaster claviger</i> , Perrier, and <i>Oreaster grayi</i> , Bell.
(3)	+	L		= <i>Oreaster Westermanni</i> , Lütken.
(4)	+	+	L		{ = <i>Pentaceros (Nidorellia) armatus</i> , Gray; <i>Oreaster armatus</i> , Müller and Troschel; <i>Gonioidiscus armatus</i> , Lütken. Incl.: <i>Gonioidiscus conifer</i> , Möbius; <i>Gonioidiscus stellata</i> , Verrill.
(5)	+	...	L		{ = <i>Gonioidiscus Michelini</i> , Perrier; <i>Oreaster armatus</i> (pars), Lütken.
(6)	+	...	L
(7)	?	...	+	L		{ = <i>Asterias obtusatus</i> , Bory de Saint Vincent; <i>Oreaster obtusatus</i> , Müller and Troschel; <i>Pentaceros obtusatus</i> , Perrier.
(8)	+	L
(9)	+	...	L
(10)	+	L
(11)	+	...	+	L		Incl.: <i>Culcita Novae Guineae</i> , Perrier (olim).
(12)	+	...	+	L		Incl.: <i>Culcita pulverulenta</i> (Val.), Perrier.
(13)	+	...	+	...	+	L		{ Incl.: <i>Randasia granulata</i> , Gray; <i>Culcita gregaria</i> , Perrier (olim).