

	LOCALITIES.	RANGE IN DEPTH: FATHOMS.	NATURE OF THE SEA-BOTTOM.	
CRYPTOZONIA.				
<i>LINCKIIDÆ.</i>				
<i>LINCKIINÆ.</i>				
<i>Ophidiaster</i> , Agassiz.				
<i>O. chinensis</i> , Perrier	Canton (China).	(1)
<i>O. cribrarius</i> , Lütken	Tonga Islands; Samoa.	(2)
* <i>O. cylindricus</i> (Lamarck), Müller and Troscchel	Mauritius; Moluccas, Fiji Islands.	} Shallow.	Coral reefs.	(3)
<i>O. duncani</i> , de Loriol	Mauritius.			...
<i>O. florida</i> , Perrier	Strait of Florida.	20	(5)
<i>O. fuscus</i> (Gray), Perrier	{ Migupou (<i>vide</i> Gray), Philippine Islands (Brit. Mus.); N.W. Australia (<i>vide</i> Studer).	}	(6)
<i>O. germani</i> , Perrier				
<i>O. gracilis</i> (Gray), Perrier	W. coast of Columbia.	(8)
<i>O. granifer</i> , Lütken	Tonga Islands.	(9)
<i>O. guildingii</i> , Gray	St. Thomas (Antilles).	(10)
* <i>O. helicostichus</i> , Sladen	Torres Strait.	6	Coral mud.	(11)
<i>O. hemprichii</i> , Müller and Troscchel	Red Sea.	(12)
<i>O. lessonæ</i> , Gasco	Capri; Mediterranean.	82	(13)
* <i>O. ophidianus</i> (Lamarck), Agassiz {	Mediterranean; Azores; Canary Islands; Cape Verde Islands.	} 5-450	Volcanic mud.	(14)
<i>O. perrieri</i> , de Loriol	Mauritius.]			
<i>O. purpureus</i> , Perrier	Mauritius; Seychelles.	(16)
<i>O. pusillus</i> , Müller and Troscchel {	Flores; Amboina; Philippine Islands; New Caledonia.	}	(17)
<i>O. pustulatus</i> (von Martens), Perrier	Amboina; Flores; Mauritius (<i>vide</i> Studer).			