

	LOCALITIES.	RANGE IN DEPTH: FATHOMS.	NATURE OF THE SEA-BOTTOM.	
CRYPTOZONIA.				
Family <i>STICHASTERIDÆ</i> , Perrier.				
<i>Stichaster</i> , Müller and Troschel.				
* <i>S. albulus</i> (Stimpson), Verrill	Greenland; northern coasts of America; Iceland; Spitzbergen.	3-192	Rocks and Hardground	(1)
<i>S. arcticus</i> , Danielssen and Koren				Off the Lofoten Islands.
* <i>S. aurantiacus</i> (Meyen), Verrill.	Peru; Chili.	Shallow.	Rocks.	(3)
<i>S. australis</i> (Verrill), Perrier	Auckland (New Zealand).	(4)
* <i>S. felipes</i> , Sladen	Off the Cape of Good Hope.	150	Green sand.	(5)
<i>S. nutrix</i> , Studer	Island of South Georgia.	(6)
* <i>S. polygrammus</i> , Sladen	W. coast of South America.	245	Blue mud.	(7)
* <i>S. polyplax</i> , Müller and Troschel, sp.	S. and S.E. Australia; Tasmania.	38-40	Sand and shells.	(8)
<i>S. roseus</i> (O. F. Müller), Sars				Coasts of Britain and Norway.
<i>S. talismani</i> , Perrier	Canary Islands; Azores.	517-782	(10)
<i>Neomorphaster</i> , Sladen.				
* <i>N. eustichus</i> , Sladen	Off the Azores.	900-1000	Pteropod ooze.	(11)
<i>Tarsaster</i> , Sladen.				
* <i>T. stoichodes</i> , Sladen	N. of Admiralty Island.	150	Coral mud.	(12)
Family <i>SOLASTERIDÆ</i> , Perrier.				
Subfamily <i>SOLASTERINÆ</i> , Sladen.				
<i>Crossaster</i> , Müller and Troschel.				
<i>C. affinis</i> , Brandt, sp.	Between Norway, Spitzbergen, and Greenland; (!) Bering Strait.	70-634	Clay (coarse, sandy, Blue); stones.	(13)