

	LOCALITIES.	RANGE IN DEPTH: FATHOMS.	NATURE OF THE SEA-BOTTOM.	
CRYPTOZONIA.				
<i>ECHINASTERIDÆ.</i>				
<i>ECHINASTERINÆ.</i>				
<i>Cribrella</i> (Agassiz), Forbes.				
* <i>C. oculata</i> (Linck), Forbes.	Off Scandinavian and British coasts; Greenland; Labrador; coast of N. America; Spitzbergen; Nova Zembla; in the White Sea; northern coast of Asia (<i>vide</i> Brandt).	Shallow water to 1350	Clay (sandy, Blue and Biloculina); Hard ground; Blue mud.	(1)
<i>C. oculata</i> , var. <i>abyssicola</i> , Norman	North Britain and Faerøe Channel.	516-555	(2)
* <i>C. ornata</i> , Perrier...	Cape of Good Hope; New Zealand; Campbell Island.	Shallow to 20	(3)
<i>C. pagenstecheri</i> , Studer	Island of South Georgia.	(1) less than 14	(4)
* <i>C. præstans</i> , Sladen	Between Marion Island and Kerguelen Island.	210	Hard ground (gravel, shells).	(5)
<i>C. sexradiata</i> , Perrier	Barbados; Gulf of Mexico.	101-321	Broken shells and corals.	(6)
* <i>C. simplex</i> , Sladen	Inaccessible Island; Nightingale Island; Marion Island; Prince Edward Island; between Marion Island and Kerguelen.	50-310	Hard ground (gravel, shells); Volcanic sand.	(7)
* <i>C. simplex</i> , var. <i>granulosa</i> , Sladen	Off Kerguelen Island.	10-50	Volcanic mud.	(8)
* <i>C. sufflata</i> , Sladen	S. of Kermadec Island.	520	Volcanic mud.	(9)
<i>Perknaster</i> , Sladen.				
* <i>P. densus</i> , Sladen	Off Kerguelen Island.	127	Volcanic mud.	(10)
* <i>P. fuscus</i> , Sladen	Off Kerguelen Island; off Heard Island.	25-75	Volcanic mud.	(11)
<i>Echinaster</i> , Müller and Troschel.				
<i>E. brasiliensis</i> , Müller and Troschel	Coast of Brazil.	(12)