

	GEOGRAPHICAL AREAS.							BATHYMETRICAL ZONES.			BOTTOM TEMPERATURE: FAHRENHEIT.	SYNONYMA AND REMARKS.
	I. North Atlantic.	II. South Atlantic.	III. Indian Ocean.	IV. Southern Ocean.	V. Eastern Archipelago.	VI. North Pacific.	VII. South Pacific.	L. Littoral, 0-150 Fathoms.	O. Continental, 150-500 Fathoms.	A. Abyssal, greater than 500 Fathoms.		
(3)	L
(3)	L
(6)	L
(4)	L	Incl.: <i>Echinaster affinis</i> , Perrier.
(6)	+	L	Incl.: <i>Echinaster Clouei</i> (Val.), Perrier.
(6)	L
(7)	+	L	C	...	46°-5-61°-75
(8)	+	...	+	L	{ = <i>Othilia purpurea</i> , Gray. Incl.: <i>Echinaster fallax</i> , Müller and Troschel; <i>Cribella fallax</i> , Dujardin and Hupé; <i>Othilia lusonica</i> , Gray.
(9)	L
(10)	+	L	41°-1
(11)	+	L	{ = <i>Asterias sentus</i> , Say. Incl.: <i>Othilia aculeata</i> , Gray; <i>Echinaster spinosus</i> (pars), Müller and Troschel; <i>Othilia spinosa</i> , Agassiz.
(12)	+	L	{ = <i>Pentadactylosaster reticulatus brevis</i> , Linck; (1) <i>Asterias seposita</i> , Lamarck; <i>Asterias</i> ("Pentastóries") <i>seposita</i> , de Blainville; <i>Stellonia seposita</i> , Nardo; <i>Rhopia seposita</i> , Gray; <i>Rhopia mediterranea</i> , Gray; <i>Cribella seposita</i> , Dujardin, and Hupé. Incl.: <i>Asterias sanguinolenta</i> , Retzius; <i>Asterias sagena</i> , Retzius.
(13)	+	L
(14)	+	+	L	{ = <i>Pentadactylosaster spinosus regularis</i> , Linck; <i>Asterias spinosa</i> , Retzius; <i>Othilia spinosa</i> , Gray. Incl.: <i>Asterias echinophora</i> , Lamarck; <i>Echinaster</i> (<i>Othilia</i>) <i>crassispina</i> , Verrill; <i>Echinaster crassispinus</i> , Lütken; <i>Echinaster echinophorus</i> , Perrier.