

	LOCALITIES.	RANGE IN DEPTH: FATHOMS.	NATURE OF THE SEA-BOTTOM.	
CRYPTOZONIA				
<i>ASTERIIDÆ.</i>				
<i>Asterias</i> , Linné.				
Subgen. <i>Stolasterias</i> , Sladen.				
* <i>A. (S.) calamaria</i> , Gray, sp.	Mauritius; S. and S.E. Australia; Tasmania; New Zealand.	0-38	(1)
.....				
* <i>A. (S.) eustyla</i> , Sladen	Off Nightingale Island (Tristan da Cunha group).	100-150	(2)
* <i>A. (S.) gemmifera</i> (Val., M.S.), Perrier, sp.	Chili; Fiji Islands.	...	Coral reefs.	(3)
.....				
* <i>A. (S.) glacialis</i> , O. F. Müller, sp.	Finmark; Iceland; European coasts; Mediterranean; Azores; Canary Islands; Cape Verde Islands.	0-66	(4)
.....				
* <i>A. (S.) stichantha</i> , Sladen	S. of Japan.	345	Green mud.	(5)
.....				
* <i>A. (S.) tenuispina</i> , Lamarck, sp.	Mediterranean; Madeira; Azores; Canary Islands; Cape Verde Islands; Bermuda; Abrolhos reefs.	0-174	(6)
.....				
* <i>A. (S.) volsellata</i> , Sladen	Off Zebu (Philippine Islands).	95	(7)
.....				
<i>Asterias</i> (not classified in subgenera).				
<i>A. acervata</i> , Stimpson	Bering Strait.	5-15	Clean gravelly bottom.	(8)
<i>A. acutispina</i> , Stimpson	Island of Ousima.	(9)
<i>A. equalis</i> , Stimpson	Monterey (California).	(10)
<i>A. alba</i> , Bell	Sandy Point (Strait of Magellan).	7-10	Dead acorn shells.	(11)
<i>A. angulosa</i> , Perrier	Alligator reefs; Strait of Florida.	85-110	(12)