

	LOCALITIES.	RANGE IN DEPTH : FATHOMS.	NATURE OF THE SEA-BOTTOM.	
CRYPTOZONIA.				
<i>BRISINGIDÆ.</i>				
<i>Brisinga</i> , Asbjørnsen.				
<i>B. mediterranea</i> , Perrier	{ Between Marseilles and Corsica.	{ 301-1454	(1)
* <i>B. membranacea</i> , Sladen	{ Between Marion Island and the Crozet Islands; and W. of the Crozet Islands.	{ 1375-1600	{ Globigerina ooze; Diatom ooze.	(2)
* <i>B. verticillata</i> , Sladen	{ E. of North America; off the coast of New Jersey.	{ 1350	Blue mud.	(3)
<i>Freyella</i> , Perrier.				
<i>F. americana</i> , Verrill, sp.	Off Nova Scotia.	175	(4)
* <i>F. attenuata</i> , Sladen	{ E. of the Mariana or Ladrone Islands.	{ 2300	Radiolarian ooze.	(5)
* <i>F. benthophila</i> , Sladen	39° 41' 0" S., 131° 23' 0" W.	2550	Red clay.	(6)
* <i>F. bracteata</i> , Sladen	{ E. of North America; S. of Nova Scotia.	{ 1250-1350	Blue mud.	(7)
* <i>F. dimorpha</i> , Sladen	Off Torres Strait, Pacific side.	1400	Globigerina ooze.	(8)
* <i>F. echinata</i> , Sladen	{ Between Celebes and Mindanao; W. of the Island of Luzon; between the Pelew Islands and New Guinea.	{ 1050-2150	{ Blue mud; Globigerina ooze.	(9)
<i>F. edwardsi</i> , Perrier	{ "Travaillour" dredgings (1 Mediterranean.)	{	(10)
<i>F. elegans</i> , Verrill, sp.	{ Between Cape Hatteras and Nova Scotia.	{ 606-2021	(11)
* <i>F. fragilissima</i> , Sladen	{ Between Marion Island and the Crozet Islands; near the Antarctic circle.	{ 1375-1975	{ Globigerina ooze; Diatom ooze.	(12)
* <i>F. heroina</i> , Sladen	{ Mid-north Pacific, 35° 22' 0" N., 169° 53' 0" E.	{ 2900	Red clay.	(13)
* <i>F. pennata</i> , Sladen	S. of Japan.	1875	Blue mud.	(14)