

	Tribe.	Family.	Genus.	Brephalos.
NORMALIA (continued.)	Monocarpidea,	THALASSOCARIDÆ,	{ <i>Thalassocaris.</i> <i>Diaphoropsis.</i> <i>Kyptocaris.</i>	
		ATYIDÆ,	{ <i>Atya,</i> <i>Caridina.</i>	Zoea.
		PONTONIIDÆ, ¹	{ <i>Pontonia.</i> (<i>Typton</i>).	
		CARIOYPHIDÆ, ¹	{ <i>Caricyphus.</i> <i>Rhomaleocaris.</i> <i>Anebocaris.</i>	
		ACANTHEPHYRIDÆ, ¹	{ <i>Bentheocaris.</i> <i>Acanthephyra.</i> <i>Systellaspis.</i> <i>Oplophorus.</i> <i>Campylonotus.</i>	
		PALÆMONIDÆ, ¹	{ <i>Palæmon,</i> <i>Palæmonella.</i> <i>Bithynis.</i> <i>Brachycarpus.</i>	Zoea.
		NEMATOCARCINIDÆ, ¹	{ <i>Nematocarcinus.</i> <i>Stochasmus.</i>	
		TROPIOCARIDÆ, ¹	{ <i>Notostomus.</i> <i>Tropiocaris.</i> <i>Hymenodora,</i>	Megalopa.
		STYLODACTYLIDÆ,	<i>Stylodactylus.</i>	
		PASIPHÆIDÆ,	{ <i>Leptochela.</i> <i>Pasiphæa.</i> <i>Orphania.</i>	
	OODEOPIDÆ,	<i>Oodeopus.</i>		
	Haplopodea,	HECTARTHROPIDÆ,	{ <i>Proctetes.</i> <i>Icotopus.</i> <i>Hectarthropus.</i> <i>Eretmocaris.</i> <i>Amphion.</i>	

Tribe Crangonidea.

This tribe contains those normal Phyllobranchiata in which the first pair of pereopoda is subchelate, and the second pair is feeble or imperfect, the carpos being uniarticulate.

Family CRANGONIDÆ.

Carapace dorsally flattened and laterally compressed. Rostrum dorsally flattened; frontal margin laterally extended; posterior margin laterally overlapped by the first somite of the pleon. Pleon laterally compressed posteriorly. Ophthalmopoda short and

¹ I have formed these families since the descriptive part was printed. For their definition see p. 927.